arXiv:math/0506350v1 [math.DG] 17 Jun 2005

RIGIDITY OF AMALGAMATED PRODUCT IN NEGATIVE CURVATURE

GÉRARD BESSON, GILLES COURTOIS, SYLVAIN GALLOT

Abstract. Let Γ be the fundamental group of a compact riemannian manifold X of sectional curvature $K \leq -1$ and dimension $n \geq 3$. We suppose that $\Gamma = A *_C B$ is the free product of its subgroups A and B over the amalgamated subgroup C. We prove that the critical exponent $\delta(C)$ of C satisfies $\delta(C) \geq n-2$. The equality happens if and only if there exist an embedded compact hypersurface $Y \subset X$, totally geodesic, of constant sectional curvature -1, whose fundamental group is C and which separates X in two connected components whose fundamental groups are A and B. Similar results hold if Γ is an HNN extension, or more generally if Γ acts on a simplicial tree without fixed point.

1. Introduction

In [15], Y. Shalom proved the following theorem which says that for every lattice Γ in the hyperbolic space and for any decomposition of Γ as an amalgamated product $\Gamma = A *_C B$, the group C has to be "big". In order to measure how "big" C is, let us define the critical exponent of a discrete group C acting on a Cartan Hadamard manifold by

$$\delta(C) = inf\{s > 0 \quad | \quad \Sigma_{\gamma\in\Gamma} e^{-sd(\gamma x, x)} < +\infty\}.$$

Theorem 1.1. *Let Γ be a lattice in $PO(n,1)$. Assume that Γ is an amalgamated product of its subgroups A and B over C. Then, the critical exponent $\delta(C)$ of C satisfy $\delta(C) \geq n-2$.*

An example is given by any n-dimensional hyperbolic manifold X which contains a compact separating connected totally geodesic hypersurface Y. The Van Kampen theorem then says that the fundamental group Γ of X is isomorphic to the free product of the fundamental groups of the two halves of $X - Y$ amalgamated over the fundamental group C of the incompressible hypersurface Y. Such examples do exist in dimension 3 thanks to the W.Thurston's hyperbolization theorem. In any dimension, A. Lubotsky showed that any standard arithmetic lattice of $PO(n,1)$ has a finite cover whose fundamental group is an amalgamated product, cf. [11]. In fact, A. Lubotsky proved that any standard arithmetic lattice Γ has a finite index subgroup Γ_0 which is mapped onto a nonabelian free group. A nonabelian free group can be written in infinitely many ways as an amalgamated product, so one get infinitely many decomposition of Γ_0 as an amalgamated

---

Date: February 1, 2008.

product by pulling back the amalgamated decomposition of the nonabelian free group.

In these cases there is equality in theorem 1.1, ie. $\delta(C) = n-2$ where C is the fundamental group of Y, and Y. Shalom suggested in [15] that the equality case in the theorem 1.1 happens only in that case.

The aim of this paper is to show that the theorem 1.1 still holds when Γ is the fundamental group of a compact riemannian manifold of variable sectional curvature less than or equal to -1, and characterize the equality case.

Theorem 1.2. *Let X be an n-dimensional compact riemanniann manifold of sectional curvature $K \leq -1$. We assume that the fundamental group Γ of X is an amalgamated product of its subgroups A and B over C and that neither A nor B equals Γ. Then, the critical exponent $\delta(C)$ of C satisfy $\delta(C) \geq n-2$. Equality $\delta(C) = n-2$ happens if and only if C cocompactly preserves a totally geodesic isometrically embedded copy $\mathbb{H}^{n-1}$ of the hyperbolic space of dimension $n-1$. Moreover, in the equality case, the hypersurface $Y^{n-1} := \mathbb{H}^{n-1}/C$ is embedded in X and separates X in two connected components whose fundamental groups are respectively A and B.*

Remark 1.3. *(i) By the assumption on A or B not being equal to Γ we exclude the trivial decomposition $\Gamma = \Gamma *_C C$ where $A = \Gamma$ and $C = B$ can be an arbitrary subgroup of Γ, for example any cyclic subgroup, in which case the conclusion of theorem 1.2 fails. Also note that because of this assumption on A and B, we have $A \neq C$ and $B \neq C$.*

(ii) Let us recall that standard arithmetic lattices in $PO(n,1)$ have finite index subgroup with infinitely many non equivalent decompositions as amalgamated products, cf. [11]. *In fact, among these decompositions, all but finitely many of them are such that $\delta(C) > n-2$. Indeed, by theorem 1.2, if $\delta(C) = n-2$ then C is the fundamental group of an embedded totally geodesic hypersurface in X, but there are only finitely many totally geodesic hypersurfaces by* [21].

When a group is an amalgamated product, it acts on a simplicial tree without fixed point and theorem 1.1 is a particular case of the

Theorem 1.4. *(*[15]*, theorem 1.6). Let $\Gamma \subset SO(n,1)$, $n \geq 3$, be a lattice Suppose Γ acts on a simplicial tree T without fixed vertex. Then there is an edge of T whose stabilizer C satisfies $\delta(C) \geq n-2$.*

In the case Γ is cocompact, the conclusion of theorem 1.4 holds for the stabilizer of any edge which separates the tree T in two unbounded components, and the proof of this is exactly the same as the proof of theorem 1.2. In particular, when the action of Γ on T is minimal, (ie. there is no proper subtree of T invariant by Γ), the conclusion of theorem 1.4 holds for every edge of T, in the variable curvature setting, and we are able to handle the equality case.

Theorem 1.5. *Let Γ be the fundamental group of an n-dimensional compact riemannian manifold X of sectional curvature less than or equal to -1. Suppose Γ acts minimally on a simplicial tree T without fixed point. Then, the stabilizer C of every edge of T satisfies $\delta(C) \geq n-2$. The equality $\delta(C) = n-2$ happens if and only if there exist a compact totally geodesic hypersurface $Y \subset X$ with fundamental group $\pi_1(Y) = C$. Moreover, in that case, Y with its induced metric has constant sectional curvature -1.*

Another interesting case contained in theorem 1.5 is the case of HNN extension. Let us recall the definition of an HNN extension. Let A and C be groups and $f_1 : C \to A$, $f_2 : C \to A$ two injective morphisms of C into A. The HNN extension $A*_C$ is the group generated by A and an element t with the relations $tf_1(\gamma)t^{-1} = f_2(\gamma)$. For example, let X be a compact manifold containing a non separating compact incompressible hypersurface $Y \subset X$. Let A be the fundamental group of the manifold with boundary $X-Y$ obtained by cuting X along Y and let C be the fundamental group of Y. The boundary of $X-Y$ consists in two connected components $Y_1 \subset X-Y$ and $Y_2 \subset X-Y$ homeomorphic to Y. By the incompressibility assumption, these inclusions give rise to two embeddings of C into A, and the fundamental group of X is the associated HNN extension $A*_C$.

Theorem 1.6. *Let Γ be the fundamental group of a compact riemannian manifold X of dimension n and sectional curvature less than or equal to -1. Suppose that $\Gamma = A*_C$ where A is a proper subgroup of Γ. Then, we have $\delta(C) \geq n-2$ and equality $\delta(C) = n-2$ if and only if there exist a non separating compact totally geodesic hypersurface $Y \subset X$ with fundamental group $\pi_1(Y) = C$. Moreover, in that case, Y with its induced metric is of constant sectional curvature -1, and the HNN decomposition arising from Y is the one we started with.*

Let us summarize the ideas of the proof of theorem 1.2. We work on $\tilde{X}/C$. The amalgamation assumption provides an essential hypersurface Z in $\tilde{X}/C$, namely Z is homologically non trivial in $\tilde{X}/C$. The volume of all hypersurfaces homologous to Z is bounded below by a positive constant because their systole are bounded away from zero. We then construct a smooth map $F : \tilde{X}/C \to \tilde{X}/C$, homotopic to the identity which contracts the volume of all compact hypersurfaces Y by the factor $\left(\frac{\delta(C)}{n-2}\right)^{n-1}$, namely $vol_{n-1}F(Y) \leq \left(\frac{\delta(C)}{n-2}\right)^{n-1} vol_{n-1}Y$. This contracting property together with the lower bound of the volume of hypersurfaces in the homology class of Z gives the inequality $\delta(C) \geq n-2$. This map is different from the map constructed in [3], in particular it can be defined under the single condition that the limit set of C is not reduced to one point. Moreover, its derivative has an upper bound depending only on the critical exponent of C.

The equality case goes as follows. When $\delta(C) = n-2$, the map $F : \tilde{X}/C \to \tilde{X}/C$ contracts the $(n-1)$-dimensional volumes, ie. $|Jac_{n-1}F| \leq 1$.

This contracting property is infinitesimally rigid in the following sense. Let us consider a lift $\tilde{F}$ of F. If $|Jac_{n-1}\tilde{F}(x)| = 1$ at some point $x \in \tilde{X}$, then $\tilde{F}(x) = x$, there exists a tangent hyperplane $E \subset T_x\tilde{X}$ such that $D\tilde{F}(x)$ is the orthogonal projector of $T_x\tilde{X}$ onto E and the limit set Λ_C is contained in the *topological equator* $E(\infty) \subset \partial\tilde{X}$ associated to E. By topological equator $E(\infty) \subset \partial\tilde{X}$ associated to E, we mean the set of end points of those geodesic rays starting at x tangently to E.

We then prove the existence of a point $x \in \tilde{X}$ such that

$$|Jac_{n-1}\tilde{F}(x)| = 1. \tag{1.1}$$

If there would exist a minimizing cycle in the homology class of Z in $\tilde{X}/C$, any point of such a cycle would satisfy (1.1). As no such minimizing cycle a priori exists because of non compactness of $\tilde{X}/C$, we prove instead the existence of a L^2 harmonic $(n-1)$-form dual to Z, which is enough to prove existence of a point x such that (1.1) holds.

At this stage of the proof, there is a big difference between the constant curvature case and the variable curvature case.

In the constant curvature case, any topological equator bounds a totally geodesic hyperbolic hypersurface $\mathbb{H}^{n-1}$, and therefore, as the group C preserves $\Lambda_C \subset E(\infty) = \partial\mathbb{H}^{n-1}$, it is not hard to see that C also preserves $\mathbb{H}^{n-1}$ and acts cocompactly on it, and the hypersurface of the equality case in theorem 1.2 is $\mathbb{H}^{n-1}/C$, [2].

In the variable curvature case, we first show the existence of a C-invariant totally geodesic hypersurface $\tilde{Z}_\infty \subset \tilde{X}$ whose boundary at infinity coincides with $\Lambda(C)$, and then we show that $\tilde{Z}_\infty$ is isometric to the real hyperbolic space. We then show that $Y =: \mathbb{H}^{n-1}_{\mathbb{R}}/C$, which is compact, injects in $X = \tilde{X}/\Gamma$ and separates X in two connected components whose fundamental groups are A and B respectively.

In order to show the existence of such a totally geodesic hypersurface $\tilde{Z}_\infty$, we first prove that C is a convex cocompact group, ie. the convex hull of the limit set of C in $\tilde{X}$ has a compact quotient under the acton of C, and that the limit set of C is homeomorphic to an $(n-2)$-dimensional sphere.

The convex cocompactness property of C and the fact that the limit set $\Lambda(C)$ of C is homeomorphic to an $(n-2)$-dimensional topological sphere are the two key points in the equality case.

This compactness property then allows us to prove the existence of a mimimizing current in the homology class of the essential hypersurface $Z \subset \tilde{X}/C$. By regularity theorem this minimizing current $\tilde{Z}_\infty$ is a smooth manifold except at a singular set of codimension at least 8. By the contracting properties of our map F, $\tilde{Z}_\infty$ is fixed by F and the geometric properties of F at fixed points where the $(n-1)$-jacobian of F equals 1 allows us to prove that $\tilde{Z}_\infty$ is totally geodesic and isometric to the hyperbolic space.

Let us now briefly describe the proof of the convex cocompactness property of C in the equality case.

The group C (or a finite index subgroup of it) actually globally preserves a smooth cocompact hypersurface $\tilde{Z} \subset \tilde{X}$ which separates $\tilde{X}$ into two connected components and whose boundary $\partial\tilde{Z} \subset \partial\tilde{X}$ coincides with $\Lambda_C \subset E(\infty)$. In the case where C wouldn't be convex cocompact, we are able to find an horoball $HB(\theta_0)$ centered at some point $\theta_0 \in \Lambda_C$ in the complementary of which lies the hypersurface $\tilde{Z}$.

The contradiction then comes from the following.

Consider a sequence of points $\theta_i \in \partial\tilde{X}$ converging to θ_0 and geodesic rays α_i starting from the point $x \in \tilde{X}$ at which $|Jac_{n-1}(x)| = 1$ and ending up at θ_i. These geodesic rays have to cross $\tilde{Z}$ at points z_i which are at bounded distance from the orbit Cx of x, therefore the shadows $\mathcal{O}_i$ of balls centered at these z_i enlighted from x have to contain points of Λ_C by the shadow lemma of D. Sullivan. On the other hand, we show that it is possible to choose the sequence θ_i in such a way that these shadows $\mathcal{O}_i$ don't meet Λ_C. This property $\mathcal{O}_i \cap \Lambda_C = \emptyset$ comes from a choice of θ_i such that the distance between z_i and the set H of all geodesics rays at x tangent to $E \subset T_x\tilde{X}$ tends to ∞. Intuitively, in order to chose z_i as far as possible from H, the points θ_i have to be chosen tranversally to Λ_C. This transversality condition is not well defined because the limit set Λ_C might be highly non regular. Thus, in order to prove that such a choice is possible, we argue again by contradiction. If for any choice of a sequence θ_i converging to θ_0, the distance between z_i and H stays bounded, then the Gromov distances $d(\theta_i, \theta_0)$ between θ_i and θ_0 satisfy $d(\theta_i, \Lambda_C) = o(d(\theta_i, \theta_0))$, and therefore any tangent cone of Λ_C at θ_0 would coincide with a tangent cone of $\partial\tilde{X}$ at θ_0, which is known to be topologically $\mathbb{R}^{n-1}$. But on the other hand, the existence of a point x such that $|Jac_{n-1}(x)| = 1$ and the fact that C acts uniformly quasiconformally with respect to the Gromov distance on $\partial\tilde{X}$ imply that the Alexandroff compactification of the above tangent cone of Λ_C at θ_0 is homeomorphic to Λ_C which is contained in a topological sphere S^{n-2}, leading to a contradiction.

From convex cocompactness of C and the fact that the limit set of C is a topological $(n-2)$-dimensional sphere, there is an alternative proof of the existence of a totally geodesic C-invariant copy of the hyperbolic space $\mathbb{H}^{n-1}_{\mathbb{R}} \subset \tilde{X}$ which consists in observing that the topological dimension and the Hausdorff dimension of the limit set $\Lambda(C)$ are equal to $n-2$ and then use the following result of M. Bonk and B. Kleiner (which we quote in the riemannian manifold setting although it remains true for $CAT(-1)$ spaces) instead of the (simpler) minimal current argument.

Theorem 1.7. [4] *Let X be a Cartan Hadamard n-dimensional manifold whose sectional curvature satisfy $K \leq -1$, and C a convex cocompact discrete subgroup of isometries of X with limit set Λ_C. Let us assume that the topological dimension and the Hausdorff dimension (with respect to the Gromov distance on $\partial\tilde{X}$) of Λ_C coincide and are equal to an integer p. Then, C*

preseves a totally geodesic embedded copy of the real hyperbolic space $\mathbb{H}^{p+1}$*, with* $\partial\mathbb{H}^{p+1} = \Lambda_C$*.*

The authors would like to express their gratitude to Alex Lubotsky, Jean Barge, Marc Bourdon, Gilles Carron, Jean Lanne, Frédéric Paulin, Leonid Potyagailo for their interests and helpfull conversations.

2. Essential hypersurfaces

Let Γ be a discrete cocompact group of isometries of a n-dimensional Cartan-Hadamard manifold $(\tilde{X}, \tilde{g})$ whose sectional curvature satisfies $K_{\tilde{g}} \leq -1$. Let us assume that the compact manifold $X = \tilde{X}/\Gamma$ is orientable. Let us also assume that $\Gamma = A *_C B$ is an amalgamated product of its subgroups A and B over C.

We first reduce to the case where $[\Gamma : C]$ is infinite.

Namely, if $[\Gamma : C] < \infty$, then the critical exponent $\delta(C) = \delta(\Gamma) \geq n-1$, and the equality in theorem 1.2 holds.

We then can assume that $[\Gamma : C] = \infty$.

Lemma 2.1. *Let* $\Gamma = A *_C B$ *be as above, with* $[\Gamma : C] = \infty$*. If neither* A *nor* B *equals* Γ*, then* $H_{n-1}(\tilde{X}/C, \mathbb{Z}) \neq 0$*.*

Proof : The Mayer-Vietoris sequence coming from the decomposition $\Gamma = A *_C B$ writes cf. [6], Corollary 7.7,

$$H_n(\tilde{X}/C, \mathbb{Z}) \to H_n(\tilde{X}/A, \mathbb{Z}) \oplus H_n(\tilde{X}/B, \mathbb{Z}) \to H_n(\tilde{X}/\Gamma, \mathbb{Z}) \to ...$$

$$... \to H_{n-1}(\tilde{X}/C, \mathbb{Z}) \to ...$$

As $[\Gamma : C] = \infty$, $H_n(\tilde{X}/C, \mathbb{Z}) = 0$ thus, if $H_{n-1}(\tilde{X}/C, \mathbb{Z}) = 0$, we deduce from the Mayer-Vietoris sequence that $H_n(\tilde{X}/A, \mathbb{Z}) \oplus H_n(\tilde{X}/B, \mathbb{Z})$ is isomorphic to $H_n(\tilde{X}/\Gamma, \mathbb{Z})$. As $H_{n-1}(\tilde{X}/\Gamma, \mathbb{Z}) = \mathbb{Z}$, we then deduce that either $[\Gamma : A] = \infty$ and $B = \Gamma$, or $[\Gamma : B] = \infty$ and $A = \Gamma$. □

In fact in the sequel of the paper we will make use of a smooth essential hypersurface Z in $\tilde{X}/C$.

Definition 2.2. *A compact smooth orientable hypersurface* Z *of an* n*-dimensional manifold* Y *is essential in* Y *if* $i_*([Z]) \neq 0$ *where* $[Z] \in H_{n-1}(Z, \mathbb{R})$ *denotes the fundamental class of* Z *and* $i_* : H_{n-1}(Z, \mathbb{R}) \to H_{n-1}(Y, \mathbb{R})$ *the morphism induced by the inclusion* $i : Z \hookrightarrow Y$*.*

The end of this section is devoted to finding such an hypersurface Z in $\tilde{X}/C$.

Let us recall a few facts about amalgamated products and their actions on trees, following [14]. Let $\Gamma = A *_C B$ be an amalgamated products of its subgroups A and B over C. Then, Γ acts on a simplicial tree $\tilde{T}$ with a fundamental domain $T \subset \tilde{T}$ being a segment, ie. an edge joining two vertices. Let us describe this tree $\tilde{T}$. There are two orbits of vertices Γv_A and Γv_B, the stabilizer of the edge v_A (resp. v_B) being A,(resp. B). There is one orbit of edges Γe_C, the stabilizer of the edge e_C being C. The fundamental domain T can be chosen as the edge e_C joining the two vertices v_A and v_B. The set of vertices adjacent to v_A, (resp. v_B), is in one to one correspondance with A/C, (resp. B/C). Note that as neither A nor B are equal to Γ, then $[A : C] \neq 1$ and $[B : C] \neq 1$, therefore for an arbitrary point t_0 on the edge e_C we see that $\tilde{T} - t_0$ is a disjoint union of two unbounded connected components. This fact will be used later on.

Let us consider a continuous Γ-equivariant map $\tilde{f} : \tilde{X} \to T$ where T is the Bass-Serre tree associated to the amalgamation $\Gamma = A *_C B$. One regularizes $\tilde{f}$ such that it is smooth in restriction to the complementary of the inverse image of the set of vertices of T. Let t_0 a regular value of $\tilde{f}$ contained in that edge of T which is fixed by the subgroup C and define $\tilde{Z} = \tilde{f}^{-1}(t_0)$. $\tilde{Z}$ is a smooth orientable possibly not connected hypersurface in $\tilde{X}$, globally C-invariant. Let us write $Z = \tilde{Z}/C$. We will show $Z \subset \tilde{X}/C$ is compact and that one of the connected components of Z is essential.

Lemma 2.3. *$Z \subset \tilde{X}/C$ is compact.*

Proof : Let us show that for any sequence $z_n \in \tilde{Z}$, there exists a subsequence z_{n_k} and $\gamma_k \in C$ such that $\gamma_k z_{n_k}$ converges. As Γ is cocompact, there exists $g_n \in \Gamma$ such that the set $(g_n z_n)$ is relatively compact. Let $g_{n_k} z_{n_k}$ a subsequence which converges to a point $z \in \tilde{X}$. By continuity, the sequence $\tilde{f}(g_{n_k} z_{n_k})$ converges to $\tilde{f}(z)$, and by equivariance we get

$$g_{n_k} \tilde{f}(z_{n_k}) = g_{n_k} t_0 \to \tilde{f}(z)$$

when k tends to ∞. As Γ acts in a simplicial way on the tree T and transitively on the set of edges, the sequence $g_{n_k} t_0$ is stationary, ie $g_{n_k} t_0 = t'_0 = g t_0$ for k large enough. Thus $g^{-1} g_{n_k} = \gamma_k \in C$ for k large enough since it fixes t_0 and $\gamma_k z_{n_k} = g^{-1} g_{n_k} z_{n_k}$ converges to $g^{-1}(z)$ $\square$

The smooth compact hypersurface Z we constructed might be not connected. Let us write $Z = Z_1 \cup Z_2 \cup ... \cup Z_k$ where the Z_j's are the connected components of Z. Each Z_j is a compact smooth oriented hypersurface of $\tilde{X}/C$.

The aim of what follows is to prove that at least one component Z_i of Z is essential.

Lemma 2.4. *There exists $i \in [1, k]$ such that Z_i is essential in $\tilde{X}/C$.*

Proof : If there exists a Z_i which doesn't separate $\tilde{X}/C$ in two connected components, then Z_i is essential in $\tilde{X}/C$. So we can assume that every Z_j, $j = 1, ...k$, does separate $\tilde{X}/C$ in two connected components. In that case we will show that there exists a Z_i which separates $\tilde{X}/C$ in two unbounded connected components which easily implies that Z_i is essential.

Let us denote U_l, $l = 1, 2, ..., p$, the connected components of $\tilde{X}/C - \cup_{j=1}^{k} Z_j$.

Claim : at least two components U_m, $U_{m'}$ are unbounded.

Assuming the claim let us finish the proof of the lemma. For each Z_j we denote V_j, V_j' the two connected components of $\tilde{X}/C - Z_j$. Then $U_m = W_1 \cap W_2 \cap ... \cap W_k$ where for each j, $W_j = V_j$ or $W_j = V_j'$. In the same way, $U_{m'} = W_1' \cap W_2' \cap ... \cap W_k'$. As $U_m \cap U_{m'} = \emptyset$, there exists $i \in [1, k]$ such that $W_i \cap W_i' = \emptyset$, thus $U_m \subset V_i$ and $U_{m'} \subset V_i'$ or $U_m \subset V_i'$ and $U_{m'} \subset V_i$ so Z_i separates $\tilde{X}/C$ into two unbounded components. This proves the lemma.

Let us prove the claim.

We have already noticed that $T - \{t_0\}$ is the disjoint union of two unbounded connected components T_1 and T_2. As C acts on T isometrically and simplicially then $T/C - \{t_0\} = T_1/C \cup T_2/C$ is the disjoint union of two unbounded connected components. Let $\bar{f} : \tilde{X}/C \to T/C$ the quotient map of $\tilde{f}$. For each component U_i, we have

$\bar{f}(U_i) \subset T_1/C$ or $\bar{f}(U_i) \subset T_2/C$, thus we can conclude the claim because $\bar{f}$ is onto $\square$

Let $\pi : \tilde{X} \to \tilde{X}/C$ be the natural projection. For any $i = 1, 2, ..., k$, let us denote $\{\tilde{Z}_i^j\}_{j \in J}$ the set of connected components of $\tilde{Z}_i =: \pi^{-1}(Z_i)$.

For each $i \in [1, k]$, we claim that C acts transitively on the set $\{\tilde{Z}_i^j\}_{j \in J}$. Namely, let us consider $\tilde{Z}_i^j$ $\tilde{Z}_i^{j'}$, $\tilde{z} \in \tilde{Z}_i^j$, $\tilde{z}' \in \tilde{Z}_i^{j'}$, and write $z = \pi\tilde{z} \in Z_i$ and $z' = \pi\tilde{z}' \in Z_i$. Let α be a continuous path on Z_i such that $\alpha(0) = z$ and $\alpha(1) = z'$, and $\tilde{\alpha}$ the lift of α such that $\tilde{\alpha}(0) = \tilde{z}$. We have $\pi\tilde{\alpha}(1) = z'$ and $\tilde{\alpha}(1) \in \tilde{Z}_i^j$ for some j, thus, there exists $c \in C$ such that $c(\tilde{\alpha}(1)) = \tilde{z}'$ and therefore $c\tilde{Z}_i^j = \tilde{Z}_i^{j'}$.$\square$

Let us denote C_i^j the stabilizer of $\tilde{Z}_i^j$, and $Z_i^j = \tilde{Z}_i^j / C_i^j \subset \tilde{X}/C_i^j$. Let us write $p : \tilde{X}/C_i^j \to \tilde{X}/C$ the natural projection.

Lemma 2.5. *The restriction of p to Z_i^j is a diffeomorphism onto Z_i. In particular, Z_i^j is compact.*

proof : Let z and z' be two points in Z_i^j such that $p(z) = p(z')$. Let $\tilde{z}$ and $\tilde{z'}$ be lifts of z and z' in $\tilde{X}$. These two points $\tilde{z}$ and $\tilde{z'}$ which are in $\tilde{Z}_i$ actually belong to the same connected component $\tilde{Z}_i^j$ because for $j \neq j'$, $\tilde{Z}_i^{j'}/C_i^j \cap \tilde{Z}_i^j/C_i^j = \emptyset$. As $p(z) = p(z')$, there exits $c \in C$ such that $\tilde{z}' = c\tilde{z}$, thus $c \in C_i^j$, and $z = z'$, therefore the restriction of p to Z_i^j is injective. The surjectivity comes from the fact that $\pi^{-1}Z_i = \bigcup_{j \in J} \tilde{Z}_i^j$ and C acts transitively on the set $\{\tilde{Z}_i^j\}_{j \in J}$. $\square$

Let us consider the integer $i \in [1, k]$ as in lemma 2.4, ie. such that $Z_i \hookrightarrow \tilde{X}/C$ is essential, and choose $\tilde{Z}_i^l$ one component of $\pi^{-1}(Z_i)$.

After possibly replacing C_i^l by an index two subgroup, we may assume that C_i^l globally preserves each of the two connected components U_i^l and V_i^l of $\tilde{X} - \tilde{Z}_i^l$.

Lemma 2.6. *Let* i, l *and* C_i^l *be chosen as above. The compact hypersurface* $Z_i^l = \tilde{Z}_i^l / C_i^l$ *is essential in* $\tilde{X}/C_i^l$*. Moreover the two connected components* U_i^l/C_i^l *and* V_i^l/C_i^l *of* $\tilde{X}/C_i^l - Z_i^l$ *are unbounded.*

Proof : Let us consider $p : \tilde{X}/C_i^l \to \tilde{X}/C$. By lemma 2.5, the restriction of p to Z_i^l is a diffeomorphism onto Z_i, therefore Z_i^l is essential in $\tilde{X}/C_i^l$ because Z_i is essential in $\tilde{X}/C$. As C_i^l preseves U_i^l and V_i^l, Z_i^l separates $\tilde{X}/C_i^l$ into two connected components U_i^l/C_i^l and V_i^l/C_i^l. and as Z_i^l is essential in $\tilde{X}/C_i^l$, U_i^l/C_i^l and V_i^l/C_i^l are unbounded. □

In the sequel of the paper we will denote $\tilde{Z}' = \tilde{Z}_i^l$, $C' = C_i^l$ and $Z' = Z_i^l = \tilde{Z}_i^l/C_i^l$.

3. Isosystolic inequality

In this section we summarize facts and results due to M.Gromov, [9]. Let Z be a p-dimensional compact orientable manifold and $i : Z \hookrightarrow Y$ an embedding of Z into Y where Y is an aspherical space. We suppose that $i_*([Z]) \neq 0$ where $i_* : H_p(Z, \mathbb{R}) \to H_p(Y, \mathbb{R})$ is the morphism induced by the embedding $Z \hookrightarrow Y$. Let us fix a riemanniann metric g on Z. For each $z \in Z$ we consider the set $\mathcal{C}_z$ of those loops α at z such that $i \circ \alpha$ is homotopically non trivial in Y.

Let us define the systole of (Z, g, i) at the point z by

Definition 3.1. $sys_i(Z, g, z) = inf\{lengh(\alpha),\ \alpha \in \mathcal{C}_z\}$

and the systole of (Z, g, i) by

Definition 3.2. $sys_i(Z, g) = inf\{sys_i(Z, g, z)\ z \in Z\}$.

The following isosystolic inequality, due to M.Gromov says that the volume of any essential submanifold Z of an aspherical space Y relatively to any riemanniann metric on Z is universally bounded below by it's systole.

Theorem 3.3. [9] *There exists a constant* C_p *such that for each* p*-dimensional riemanniann manifold* (Z, g) *and any embedding* $Z \hookrightarrow Y$ *into an aspherical space* Y *such that* $i_*([Z]) \neq 0$ *where* $i_* : H_p(Z, \mathbb{R}) \to H_p(Y, \mathbb{R})$ *is the induced morphism in homology, then* $vol_p(Z, g) \geq C_p(sys_i(Z, g))^p$

We will apply this volume estimates to the essential hypersurface $i : Z \hookrightarrow \tilde{X}/C$ that we constructed in lemma 2.6.

The following lemma is immediate.

Lemma 3.4. *Let C be a discrete group acting on a simply connected manifold $\tilde{X}$, $\tilde{Z}$ a C-invariant hypersurface of $\tilde{X}$ and $i : Z = \tilde{Z}/C \hookrightarrow \tilde{X}/C$ the natural inclusion. Let g any riemanniann metric on Z and $\tilde{g}$ the lift of g to $\tilde{Z}$. Then, for any $z \in Z$ we have,*

$$sys_i(Z, g, z) = inf\{d_{\tilde{g}}(\tilde{z}, \gamma\tilde{z}), \gamma \in C\}$$

where $\tilde{z} \in \tilde{Z}$ is a lift of $z \in Z$ and $d_{\tilde{g}}$ is the distance induced by $\tilde{g}$ on $\tilde{Z}$.

Proof : Let $\alpha \in \mathcal{C}_z$ a loop based at $z \in Z$. As $i \circ \alpha$ is an homotopically non trivial loop at $i(z) = z$ in $\tilde{X}/C$, its lift $\widetilde{i \circ \alpha}$ at some $\tilde{z} \in \tilde{Z}$ ends up at $\gamma\tilde{z}$ for some $\gamma \in C$.□

4. VOLUME OF HYPERSURFACES IN $\tilde{X}/C$

Let $(\tilde{X}, \tilde{g})$ be a n-dimensional Cartan-Hadamard manifold whose sectional curvature satisfies $K_{\tilde{g}} \leq -1$ and C a discrete group of isometries of $(\tilde{X}, \tilde{g})$. We assume that the group C is non elementary, namely C fixes neither one nor two points in the geometric boundary $\partial\tilde{X}$ of $(\tilde{X}, \tilde{g})$.

The aim of this section is to construct a map $F : \tilde{X}/C \to \tilde{X}/C$ such that for any compact hypersurface Z of $\tilde{X}/C$, we have

$$vol_{n-1}(F(Z)) \leq \Big(\frac{\delta + 1}{n - 1}\Big)^{n-1} vol_{n-1}(Z)$$

where δ is the critical exponent of C and $vol_{n-1}(Z)$ stands for the $(n-1)$-dimensional volume of the metric on Z induced from g. For every subgroup $C' \subset C$ and any hypersurface Z' of $\tilde{X}/C'$ the lift $F' : \tilde{X}/C' \to \tilde{X}/C'$ of F will also verify

$$vol_{n-1}(F'(Z')) \leq \Big(\frac{\delta + 1}{n - 1}\Big)^{n-1} vol_{n-1}(Z').$$

In order to construct the map F we need a few prelimiraries. We consider a finite positive Borel measure μ on the boundary $\partial\tilde{X}$ whose support contains at least two points. Let us fix an origin $o \in \tilde{X}$ and denote $B(x, \theta)$ the Busemann function defined for each $x \in \tilde{X}$ and $\theta \in \partial\tilde{X}$ by

$$B(x, \theta) = \lim_{t\to\infty} dist(x, c(t)) - t$$

where $c(t)$ is the geodesic ray such that $c(0) = o$ and $c(+\infty) = \theta$.

Let $\mathcal{D}_\mu : \tilde{X} \to \mathbb{R}$ the function defined by

$$\mathcal{D}_\mu(y) = \int_{\partial\tilde{X}} e^{B(y,\theta)} d\mu(\theta) \tag{4.1}$$

A computation shows that

$$Dd\mathcal{D}_\mu(y) = \int_{\partial\tilde{X}} (DdB(y, \theta) + DB(y, \theta) \otimes DB(y, \theta))e^{B(y,\theta)} d\mu(\theta). \tag{4.2}$$

When $K_{\tilde{g}} \leq -1$ the Rauch comparison theorem says that for every $y \in \tilde{X}$, and $\theta \in \partial\tilde{X}$,

$$DdB(y,\theta) + DB(y,\theta) \otimes DB(y,\theta) \geq \tilde{g}. \tag{4.3}$$

We then get

$$Dd\mathcal{D}_\mu(y) \geq \mathcal{D}_\mu(y)\tilde{g}, \tag{4.4}$$

thus $Dd\mathcal{D}_\mu(y)$ is positive definite and $\mathcal{D}_\mu$ is strictly convex.

Lemma 4.1. *We have* $\lim_{y_k \to \partial\tilde{X}} \mathcal{D}_\mu(y) = +\infty$.

proof : Let $y_k \in \tilde{X}$ a sequence such that

$$\lim_{k\to\infty} y_k = \theta_0 \in \partial\tilde{X}. \tag{4.5}$$

As $supp(\mu) \cap (\partial\tilde{X} - \{\theta_0\}) \neq \emptyset$, there exists a compact subset $K \subset \partial\tilde{X} - \{\theta_0\}$ such that $\mu(K) > 0$ thus,

$$\int_{\partial\tilde{X}} e^{B(y_k,\theta)} d\mu \geq \int_K e^{B(y_k,\theta)} d\mu \to +\infty. \tag{4.6}$$

□

Corollary 4.2. *Let μ a finite borel measure on $\partial\tilde{X}$ whose support contains at least two points. The function $\mathcal{D}_\mu$ has a unique minimum. This minimum will be denoted by $\mathcal{C}(\mu)$.*

Let us now consider some discrete subgroup $C \subset Isom(\tilde{X},\tilde{g})$. Recall that a family of Patterson measures $(\mu_x)_{x\in\tilde{X}}$ associated to C is a set of positive finite measures μ_x on $\partial\tilde{X}$, $x \in \tilde{X}$, such that the following holds for all $x \in \tilde{X}$, $\gamma \in C$,

$$\mu_{\gamma x} = \gamma_* \mu_x \tag{4.7}$$

$$\mu_x = e^{-\delta B(x,\theta)} \mu_o, \tag{4.8}$$

where $o \in \tilde{X}$ is a fixed origin, B the Busemann function associated to o and δ the critical exponent of C.

We assume now that $supp(\mu_o)$ contains at least two points and define the map $\tilde{F} : \tilde{X} \to \tilde{X}$ for $x \in \tilde{X}$ by

$$\tilde{F}(x) = \mathcal{C}(e^{-B(x,\theta)}\mu_x). \tag{4.9}$$

Here are a few notations. For a subspace E of $T_x\tilde{X}$, we will write $Jac_E\tilde{F}(x)$ the determinant of the matrix of the restriction of $D\tilde{F}(x)$ to E with respect to orthonormal bases of E and $D\tilde{F}(x)E$. For an integer p, we denote by $Jac_p\tilde{F}(x)$ the supremum of $|Jac_E\tilde{F}(x)|$ as E runs through the set of p-dimensional subspaces of $T_x\tilde{X}$.

Lemma 4.3. *The map $\tilde{F}$ is smooth, homotopic to the Identity and verifies for all $x \in \tilde{X}$, $\gamma \in C$ and $p \in [2, n = dim(X)]$,*

(i) $\tilde{F}(\gamma x) = \gamma\tilde{F}(x)$

(ii) $|Jac_p\tilde{F}(x)| \leq \left(\frac{(\delta+1)}{p}\right)^p$.

Proof :
The map

$$(x,y) \rightarrow \int_{\partial\tilde{X}} e^{B(y,\theta)-B(x,\theta)}d\mu_x(\theta) = \int_{\partial\tilde{X}} e^{B(y,\theta)-(\delta+1)B(x,\theta)}d\mu_o(\theta)$$

is smooth because $y \rightarrow B(y,\theta)$ is smooth.

For all x the map $y \rightarrow \int_{\partial\tilde{X}} e^{B(y,\theta)-(\delta+1)B(x,\theta)}d\mu_o(\theta)$ is strictly convex by (4.4) and tends to infinity when y tend to $\partial\tilde{X}$ (cf. lemma 4.1), thus the unique minimum $\tilde{F}(x)$ is a smooth function. The equivariance of $\tilde{F}$ comes from the cocycle relation $B(\gamma y, \gamma\theta) - B(\gamma x, \gamma\theta) = B(y,\theta) - B(x,\theta)$.

For each $x \in \tilde{X}$ let c_x be the geodesic in $\tilde{X}$ such that $c_x(0) = x$, $c_x(1) = \tilde{F}(x)$ and which is parametrized with constant speed. The map $\tilde{F}_t : \tilde{X} \rightarrow \tilde{X}$ defined by $\tilde{F}_t(x) = c_x(t)$ is a C-equivariant homotopy between $Id_{\tilde{X}}$ and $\tilde{F}$.

It remains to prove (ii).

The point $\tilde{F}(x)$ is characterized by

$$\int_{\partial\tilde{X}} DB(\tilde{F}(x),\theta)e^{B(\tilde{F}(x),\theta)-B(x,\theta)}d\mu_x(\theta) = 0. \tag{4.10}$$

In order to simplify the notations we will write $B_{(x,\theta)}$ instead of $B(x,\theta)$ and we will denote ν_x the measure $e^{B(\tilde{F}(x),\theta)-B(x,\theta)}\mu_x$. We will also write $D\tilde{F}(u)$ instead of $D\tilde{F}(x)(u)$.

The differential of $\tilde{F}$ is characterized by the following: for $u \in T_x\tilde{X}$ and $v \in T_{\tilde{F}(x)}\tilde{X}$, one has

$$\int_{\partial\tilde{X}} [DdB_{(\tilde{F}(x),\theta)}(D\tilde{F}(u),v) + DB_{(\tilde{F}(x),\theta)}(v)DB_{(\tilde{F}(x),\theta)}(D\tilde{F}(u))]d\nu_x(\theta)$$

$$= (\delta+1)\int_{\partial\tilde{X}} DB_{(\tilde{F}(x),\theta)}(v)DB_{(x,\theta)}(u)d\nu_x(\theta). \tag{4.11}$$

We define the quadratic forms k and h for $v \in T_{\tilde{F}(x)}\tilde{X}$ by

$$k(v,v) = \int_{\partial\tilde{X}} [DdB_{(\tilde{F}(x),\theta)}(v,v) + (DB_{(\tilde{F}(x),\theta)}(v))^2]d\nu_x(\theta). \tag{4.12}$$

and

$$h(v,v)=\int_{\partial\tilde{X}} DB_{(\tilde{F}(x),\theta)}(v)^2 d\nu_x(\theta). \tag{4.13}$$

The relation (4.11) writes, for $u\in T_x\tilde{X}$ and $v\in T_{\tilde{F}(x)}\tilde{X}$:

$$k(D\tilde{F}(u),v)=(\delta+1)\int_{\partial\tilde{X}} DB_{(\tilde{F}(x),\theta)}(v)DB_{(x,\theta)}(u)d\nu_x(\theta). \tag{4.14}$$

We defines the quadratic form h' on $T_x\tilde{X}$ for $u\in T_x\tilde{X}$ by

$$h'(u,u)=\int_{\partial\tilde{X}} DB_{(x,\theta)}(u)^2 d\nu_x(\theta), \tag{4.15}$$

and one derives from (4.14)

$$|k(D\tilde{F}(x)(u),v)|\leq(\delta+1)h(v,v)^{1/2}h'(u,u)^{1/2}. \tag{4.16}$$

One now can estimate $Jac_p\tilde{F}(x)$. Let $P\subset T_x\tilde{X}$, $dimP=p$. If $D\tilde{F}(P)$ has dimension lower than p, then there is nothing to be proven. Let us assume that $dimD\tilde{F}(P)=p$. Denote by the same letters H' [resp. H and K] the selfadjoint operators (with respect to $\tilde{g}$) associated to the quadratic forms h' [resp. h, k] restricted to P [resp. $D\tilde{F}(P)$].

Let $(v_i)_{i=1}^p$ an orthonormal basis of $D\tilde{F}(P)$ which diagonalizes H and $(u_i)_{i=1}^p$ an orthonormal basis of P such that the matrix of $K\circ D\tilde{F}(x):P\to D\tilde{F}(P)$ is triangular. Then,

$$detK.|Jac_P\tilde{F}(x)|\leq(\delta+1)^p(\Pi_{i=1}^p h(v_i,v_i)^{1/2})(\Pi_{i=1}^p h'(u_i,u_i)^{1/2}) \tag{4.17}$$

thus,

$$detK.|Jac_P\tilde{F}(x)|\leq(\delta+1)^p\Big(\frac{TraceH}{p}\Big)^{p/2}\Big(\frac{TraceH'}{p}\Big)^{p/2}. \tag{4.18}$$

In these inequalities one can normalize the measures

$$\nu_x=e^{B(\tilde{F}(x),\theta)-B(x,\theta)}\mu_x$$

such that their total mass equals one, which gives

$$traceH=\Sigma_{i=1}^p h(v_i,v_i)\leq 1, \tag{4.19}$$

the last inequality coming from the fact that for all $\theta\in\partial\tilde{X}$,

$$\Sigma_{i=1}^p DB(\tilde{F}(x),\theta)(v_i)^2\leq||\nabla B(\tilde{F}(x),\theta)||^2=1 \tag{4.20}$$

and from the previous normalization.

Similarly,

$$traceH' = \Sigma_{i=1}^{p} h'(u_i, u_i) \leq 1. \tag{4.21}$$

We then obtain with (4.18)

$$detK.|Jac_P \tilde{F}(x)| \leq \Big(\frac{\delta+1}{p}\Big)^p. \tag{4.22}$$

Thanks to (4.3), we have $detK \geq 1$, so

$$|Jac_P \tilde{F}(x)| \leq \Big(\frac{\delta+1}{p}\Big)^p. \tag{4.23}$$

We get (ii) by taking the supremum in P. $\square$

As the map $\tilde{F} : \tilde{X} \to \tilde{X}$ is C-equivariant, then for every subgroup $C' \subset C$, $\tilde{F}$ gives rise to a map $F' : \tilde{X}/C' \to \tilde{X}/C'$ and so does the homotopy $\tilde{F}_t$ between $\tilde{F}$ and $Id_{\tilde{X}}$.

Corollary 4.4. *The map $F' : \tilde{X}/C' \to \tilde{X}/C'$ is homotopic to the Identity map and verifies for all $x \in \tilde{X}/C'$ and $p \in [2, n = dimX]$*

$$|JacF'_p(x)| \leq \Big(\frac{\delta+1}{p}\Big)^p.$$

Let $C \subset Isom(\tilde{X}, \tilde{g})$ as above, ie such that the support of the Patterson-Sullivan measures contains at least two points and with critical exponent δ. Let $C' \subset C$ be a subgroup.

Let us consider an compact hypersurface $Z' \subset \tilde{X}/C'$.

Denote $F'^k = F' \circ F' \circ ..... \circ F'$ the composition of F' k-times.

Let us write $g_k = (F'^k)^* g$, where g is the metric on $\tilde{X}/C'$ induced by $\tilde{g}$. The symmetric 2-tensor g_k may not be a riemanniann metric on $\tilde{X}/C'$ nor its resriction to Z', so we have to modify it. For $\epsilon > 0$, the following symmetric 2-tensor $g_{\epsilon,k}$ is a riemanniann metric on $\tilde{X}/C'$ and so is its restriction $h_{\epsilon,k}$ to Z'.

$$g_{\epsilon,k} = g_k + \epsilon^2 g. \tag{4.24}$$

Lemma 4.5. *Let $h_{\epsilon,k}$ be the restriction of $g_{\epsilon,k}$ to the hypersurface Z' and $g_{Z'}$ the restriction of g to Z'. Let $\Phi_{\epsilon,k} : Z' \to \mathbb{R}$ the density defined for all $x \in Z'$ by $dv_{h_{\epsilon,k}}(x) = \Phi_{\epsilon,k}(x) dv_{g_{Z'}}(x)$. For any sequence ϵ_k such that $\lim_{k\to\infty} \epsilon_k = 0$, there exists a sequence ϵ'_k, $\lim_{k\to\infty} \epsilon'_k = 0$, such that for all $x \in Z'$,*

$$0 < \Phi_{\epsilon'_k,k}(x) \leq |Jac_{n-1} F'^k(x)| + \epsilon_k.$$

In particular,

$$\Phi_{\epsilon'_k,k}(x) \leq \Big(\frac{\delta+1}{n-1}\Big)^{k(n-1)} + \epsilon_k$$

and

$$vol(Z', h_{\epsilon'_k,k}) \leq \Big[\Big(\frac{\delta+1}{n-1}\Big)^{k(n-1)} + \epsilon_k\Big] vol(Z', g_{Z'}).$$

Corollary 4.6. *Under the above asumptions, if $\delta < n-2$ there exists a sequence ϵ'_k such that $\lim_{k\to\infty}\epsilon'_k = 0$, and $lim_{k\to\infty} vol(Z', h_{\epsilon'_k,k}) = 0$.*

Proof of lemma 4.5 :

Let us fix k an integer. Let $x \in Z'$ and $u \in T_xZ'$. We have $g_{\epsilon,k}(u,u) = h_{\epsilon,k}(u,u) = g(DF'^k(x)(u), DF'^k(x)(u)) + \epsilon^2 g(u,u)$ thus $h_{\epsilon,k}(u,u) = g(A_{x,\epsilon}u,u)$ where $A_{x,\epsilon} \in End(T_xZ')$ is the self adjoint operator $A_x = DF'^k(x)^* \circ DF'^k(x) + \epsilon^2 Id$, with $DF'^k(x)^*$ the adjoint of $DF'^k(x) : (T_xZ', g(x)) \to (DF'^k(x)(T_xZ'), g(F'^k(x)))$.

By compactness of Z' and continuity of $A_{x,\epsilon}$, there exist ϵ'_k such that

$$\Phi_{\epsilon'_k,k}(x) = det A^{1/2}_{x,\epsilon'_k} \leq det A_{x,0} + \epsilon_k,$$

thus

$$\Phi_{\epsilon'_k,k}(x) \leq |Jac_{n-1}F'^k(x)| + \epsilon_k.$$

The lemma then follows from Corollary 4.4. □

5. Proof of Theorem 1.2

This section is devoted to the proof of the Theorem 1.2. Let Γ be a discrete cocompact group of isometries of a n-dimensional Cartan-Hadamard manifold $(\tilde{X}, \tilde{g})$ whose sectional curvature satisfies $K_{\tilde{g}} \leq -1$. We assume that $\Gamma = A *_C B$. At the end of section 2 we constructed a subgroup $C' \subset C$ and an orientable hypersurface $\tilde{Z}' \subset \tilde{X}$ such that $C'.\tilde{Z}' = \tilde{Z}'$ and $Z' = \tilde{Z}'/C'$ is compact in $\tilde{X}/C'$. Moreover Z' is essential in $\tilde{X}/C'$ ie $i_*([Z']) \neq 0$ where $i_* : H_{n-1}(Z', \mathbb{R}) \to H_{n-1}(\tilde{X}/C', \mathbb{R})$ is the morphism induced on homology groups by the inclusion $i : Z' \to \tilde{X}/C'$ and $[Z']$ the fundamental class of Z'.

5.1 Proof of the inequality

We now prove the inequality in the theorem 1.2. Let us assume that $\delta < n-2$ and derive a contradiction. Let $h_{\epsilon'_k,k}$ the sequence of metric defined on Z' in lemma 4.5, then by corollary 4.6 we have

$$\lim_{k\to\infty} vol(Z', h_{\epsilon'_k,k}) = 0. \tag{5.1}$$

We now show that the systole of the metric $h_{\epsilon'_k,k}$ on Z' is bounded below independently of k. Recall that the systole of $i : Z' \to \tilde{X}/C'$ at a point $z \in Z'$ with respect to a metric $h_{\epsilon,k}$ can be defined by

$$\mathrm{sys_i(Z', h_{\epsilon,k}, z) = inf_{\gamma\in C'} dist_{(\tilde{Z}', \tilde{h}_{\epsilon,k})}(\tilde{z}, \gamma\tilde{z})} \tag{5.2}$$

where $\tilde{z}$ is any lift of z and $\tilde{h}_{\epsilon,k}$ the lift on $\tilde{Z}'$ of $h_{\epsilon,k}$, (cf. lemma 3.4).

Let $\alpha(t)$ be a minimizing geodesic between $\tilde z$ and $\gamma\tilde z$ on $(Z',\tilde h_{\epsilon,k})$. By definition of $\tilde h_{\epsilon,k}$ we have

$$\mathrm{dist}_{(\tilde Z',\tilde h_{\epsilon,k})}(\tilde z,\gamma\tilde z)\geq l_{\tilde g}(\tilde F^k\circ\alpha) \tag{5.3}$$

where $l_{\tilde g}$ stands for the lengh with respect to $\tilde g$ on $\tilde X$.
We get

$$\mathrm{dist}_{(\tilde Z',\tilde h_{\epsilon,k})}(\tilde z,\gamma\tilde z)\geq \mathrm{dist}_{(\tilde X,\tilde g)}(\tilde F^k(\tilde z),\gamma\tilde F^k(\tilde z))\geq\rho \tag{5.4}$$

where ρ is the injectivity radius of $\tilde X/C'$.
We then have

$$\mathrm{sys}_i(Z',h_{\epsilon,k})\geq\rho, \tag{5.5}$$

and thanks to the Theorem 3.3 we obtain

$$\mathrm{vol}(Z',h_{\epsilon'_k,k})\geq C_n\rho^{n-1} \tag{5.6}$$

which contradicts (5.1).
□

5.2 Proof of the equality case
There will be several steps.

Step 1: The limit set of C is contained in a topological equator.

Step 2: The weak tangent to $\partial\tilde X$ and $\Lambda_{C'}$.

Step 3: The limit set $\Lambda_{C'}$ of C' and the limit set Λ_C of C are equal to a topological equator.

Step 4: C' and C are convex cocompact.

Step 5: C preserves a copy of the real hyperbolic space $\mathbb{H}^{n-1}_{\mathbb{R}}$ totally geodesically embedded in $\tilde X$.

Step 6: Conclusion

Step 1: The limit set Λ_C of C is contained in a topological equator.

Let $x\in\tilde X$ and $E\subset T_x\tilde X$ a codimension one subspace. For each $u\in T_x\tilde X$, $\tilde g(u,u)=1$, one considers the geodesic c_u defined by $c_u(0)=x$ and $\dot{c_u}(0)=u$. We define the equator $E(\infty)$ associated to E as the subset of $\partial\tilde X$

$$E(\infty)=\{c_u(+\infty)/u\in E\} \tag{5.7}$$

Our goal is to prove the existence of a point $x \in \tilde{X}$ and an hyperplane $E \subset T_x\tilde{X}$ such that the limit set Λ_C satisfies $\Lambda_C \subset E(\infty)$.

Recall that $C' \subset C$ globally preserves an hypersurface $\tilde{Z}'$ such that $\tilde{Z}'/C' \subset \tilde{X}/C'$ is compact and essential.

Let us also recall that we have constructed a C-equivariant map $\tilde{F} : \tilde{X} \to \tilde{X}$ such that, for all $x \in \tilde{X}$,

$$|Jac_{n-1}\tilde{F}(x)| \leq \Big(\frac{\delta+1}{n-1}\Big)^{n-1} \tag{5.8}$$

where the critical exponent δ of C satisfies $\delta = n-2$, thus

$$|Jac_{n-1}\tilde{F}(x)| \leq 1. \tag{5.9}$$

The step 1 follows from the two following Propositions.

Proposition 5.1. *Let $x \in \tilde{X}$ such that $|Jac_{n-1}\tilde{F}(x)| = 1$. Then there exists $E \subset T_x\tilde{X}$ such that the limit set Λ_C satisfies $\Lambda_C \subset E(\infty)$. Moreover, $\tilde{F}(x) = x$ and $D\tilde{F}(x)$ is the orthogonal projector onto E.*

Proposition 5.2. *There exists $x \in \tilde{X}$ such that $|Jac_{n-1}\tilde{F}(x)| = 1$.*

Proof of Proposition 5.1

Let $x \in \tilde{X}$ such that $|Jac_{n-1}\tilde{F}(x)| = 1$. By definition we have a subspace $E \subset T_x\tilde{X}$ such that $|Jac_E\tilde{F}(x)| = 1$. By (4.18) and detK ≥ 1 we have,

$$|Jac_E\tilde{F}(x)| \leq (n-1)^{n-1}\Big(\frac{traceH}{n-1}\Big)^{\frac{n-1}{2}}\Big(\frac{traceH'}{n-1}\Big)^{\frac{n-1}{2}}$$

$$\leq (n-1)^{n-1}(\frac{1}{n-1})^{n-1}. \tag{5.10}$$

In particular as $|Jac_E\tilde{F}(x)| = 1$, we have equality in the inequalities (5.10), thus, $traceH = trace(h) = 1$, and

$$H = \frac{1}{n-1}Id_{D\tilde{F}(x)(E)}. \tag{5.11}$$

Let us recall that the quadratic form h is defined by

$$h(v,v) = \int_{\partial\tilde{X}} DB(\tilde{F}(x),\theta)(v)^2 e^{B(\tilde{F}(x),\theta)-B(x,\theta)}d\mu_x(\theta)$$

where μ_x is the Patterson-Sullivan measure of C normalized by

$$\int_{\partial\tilde{X}} e^{B(\tilde{F}(x),\theta)-B(x,\theta)}d\mu_x(\theta) = 1. \tag{5.12}$$

We then have

$$1 = trace(h) = traceH = \Sigma_{i=1}^{n-1} h(v_i, v_i) =$$

$$= \int_{\partial \tilde{X}} \Sigma_{i=1}^{n-1} DB(\tilde{F}(x), \theta)(v_i)^2 e^{B(\tilde{F}(x),\theta)-B(x,\theta)} d\mu_x(\theta)$$

$$\leq \int_{\partial \tilde{X}} ||DB(\tilde{F}(x), \theta)||^2 e^{B(\tilde{F}(x),\theta)-B(x,\theta)} d\mu_x(\theta) \leq 1,$$

because $\Sigma_{i=1}^{n-1} DB(\tilde{F}(x), \theta)(v_i)^2 \leq ||DB(\tilde{F}(x), \theta)||^2 = 1$
for all $\theta \in \partial \tilde{X}$.

Therefore for μ_x-almost all $\theta \in supp(e^{B(\tilde{F}(x),\theta)-B(x,\theta)} d\mu_x(\theta)) = supp(\mu_x)$, we have

$$\Sigma_{i=1}^{n-1} DB(\tilde{F}(x), \theta)(v_i)^2 = ||DB(\tilde{F}(x), \theta)||^2 = 1. \tag{5.13}$$

In (5.12), $\Sigma_{i=1}^{n-1} DB(\tilde{F}(x), \theta)(v_i)^2$ represents the square of the norm of the projection of $\nabla B(\tilde{F}(x), \theta)$ on E.

By continuity of $B(x, \theta)$ in θ one then gets $\Lambda_C = supp(\mu_x) \subset E(\infty)$.

Let us now prove that $\tilde{F}(x) = x$. When $Jac\tilde{F}_E(x) = 1$, we have equality in the Cauchy-Schwarz inequality (4.16), therefore for each $i = 1, ...n-1$ and $\theta \in \Lambda_C$ we get $DB(\tilde{F}(x), \theta)(v_i) = DB(x, \theta)(u_i)$. Therefore we deduces from (5.13) that $\nabla B(x, \theta) = \Sigma_{i=1}^{i=n-1} DB(x, \theta)(u_i) u_i$, which imply with (4.10) that

$$\int_{\partial \tilde{X}} DB(x, \theta) e^{B(\tilde{F}(x),\theta)-B(x,\theta)} d\mu_x(\theta) = 0. \tag{5.14}$$

On the other hand, as $\int_{\partial \tilde{X}} DB(\tilde{F}(x), \theta) e^{B(\tilde{F}(x),\theta)-B(x,\theta)} d\mu_x(\theta) = 0$ and $H = \frac{1}{n-1} Id_{D\tilde{F}(x)(E)}$, the support of μ_x cannot be just a pair of points, therefore the barycenter of the measure $e^{B(\tilde{F}(x),\theta)-B(x,\theta)} \mu_x$ defined in [2] is well defined and characterized as the point $z \in \tilde{X}$ such that

$$\int_{\partial \tilde{X}} DB(z, \theta) e^{B(\tilde{F}(x),\theta)-B(x,\theta)} d\mu_x(\theta) = 0,$$

thus (5.14) and (4.10) imply $x = \tilde{F}(x)$. □

Proof of Proposition 5.2 :

If we knew that there exists a minimizing hypersurface Z_0 in the homology class of Z, then every points $x \in Z_0$ would verify $|Jac_{n-1}\tilde{F}(x)| = 1$. We unfortunately don't know if there exists such a minimizing hypersurface nor a minimizing current in the homology class of Z. Instead we will consider an $L^2(\tilde{X}/C')$ harmonic $(n-1)$-form dual to the homology class of Z.

We need the following lemmas in order to prove the existence of such a dual form.

Let $\lambda_1(\tilde{X}/C')$ be the bottom of the spectrum of the Laplacian on $(\tilde{X}, \tilde{g})$, ie.

$$\lambda_1(\tilde{X}/C') = inf_{u\in C_0^\infty(\tilde{X}/C')}\{\frac{\int_{\tilde{X}/C'}|du|^2}{\int_{\tilde{X}/C'}u^2}\}. \tag{5.15}$$

Lemma 5.3. *Let $C \subset Isom(\tilde{X}, \tilde{g})$ a discrete group of isometries with critical exponent $\delta = n-2$ where $(\tilde{X}, \tilde{g})$ is an n-dimensional Cartan-Hadamard manifold of sectional curvature $K_{\tilde{g}} \leq -1$. Then for any subgroup $C' \subset C$ we have $\lambda_1(\tilde{X}/C') \geq n-2$.*

Proof :

Thanks to a theorem of Barta, cf.[17] Theorem 2.1, the lemma boils down to finding a positive function $c : \tilde{X}/C' \to \mathbb{R}_+$ such that $\Delta c(x) \geq (n-2)c(x)$. Here, the laplacian Δ is the positive operator ie. $\Delta c = -traceDdc$. We consider the smooth function $\tilde{c} : \tilde{X} \to \mathbb{R}_+$ defined by $\tilde{c}(x) = \mu_x(\partial\tilde{X})$ where $\{\mu_x\}_{x\in\tilde{X}}$ is a family of Patterson-Sullivan measure of C. The function $\tilde{c}$ is C-equivariant therefore it defines a map $c : \tilde{X}/C' \to \mathbb{R}_+$ for any subgroup $C' \subset C$. Let us show

$$\Delta c(x) \geq \delta(n-1-\delta)c(x) = n-2. \tag{5.16}$$

We have

$$\tilde{c}(x) = \int_{\partial\tilde{X}} e^{-\delta B(x,\theta)} d\mu_o(\theta)$$

therefore

$$\Delta\tilde{c}(x) = \int_{\partial\tilde{X}} [-\delta\Delta B(x,\theta) - \delta^2] d\mu_x(\theta).$$

The sectional curvature $K_{\tilde{g}}$ of $(\tilde{X}, \tilde{g})$ satisfies $K_{\tilde{g}} \leq -1$ we thus have $-\Delta B(x,\theta) \geq n-1$ and as $\delta = n-2$ we get

$$\Delta\tilde{c}(x) \geq [\delta(n-1) - \delta^2]\tilde{c}(x) = (n-2)\tilde{c}(x).$$

□

The following lemma is due to G.Carron and E.Pedon, [8]. For a complete riemannian manifold Y, we denote $H_c^1(Y, \mathbb{R})$ the first cohomology group generated by diferential forms with compact support.

Lemma 5.4 ([8], Lemme 5.1)**.** *Let Y be a complete riemannian manifold all ends of whose having infinite volume and such that $\lambda_1(Y) > 0$, then the natural morphism*

$$H_c^1(Y, \mathbb{R}) \to H_{L^2}^1(Y, \mathbb{R})$$

is injective. In particular any $\alpha \in H_c^1(Y, \mathbb{R})$ admits a representative $\bar{\alpha}$ which is in $L^2(Y, \mathbb{R})$.

Corollary 5.5. *Let C' be as above and assume that there exists a compact essential hypersurface $Z' \subset \tilde{X}/C'$. Then there exists an harmonic $n-1$-form ω in $L^2(\tilde{X}/C')$ such that $\int_{Z'} \omega \neq 0$.*

Proof :

Let $\alpha \in H^1_c(\tilde{X}/C', \mathbb{R})$ a Poincaré dual of $[Z'] \in H_{n-1}(\tilde{X}/C', \mathbb{R})$. By definition of α, for any $\beta \in H^{n-1}(\tilde{X}/C', \mathbb{R})$, one has

$$\int_{Z'} \beta = \int_{\tilde{X}/C'} \beta \wedge \alpha, \tag{5.17}$$

([5] p.51, note that $\tilde{X}/C'$ has a " finite good cover").

After Lemma 5.4 , α admits a non trivial harmonic representative $\bar{\alpha}$ in $L^2(\tilde{X}/C')$. (In order to apply the Lemma 5.4, one has to check that all ends of $\tilde{X}/C'$ have infinite volume, ie for a compact $K \subset \tilde{X}/C'$ each unbounded connected component of $\tilde{X}/C' - K$ has infinite volume: this comes from the fact that the injectivity radius of $\tilde{X}/C'$ is bounded below by the injectivity radius of $X = \tilde{X}/\Gamma$ and the sectional curvature bounded above by -1.) The $(n-1)$-harmonic form $\omega = *\bar{\alpha}$, where $*$ is the Hodge operator, is in $L^2(\tilde{X}/C')$ and verifies after (5.17)

$$\int_{Z'} \omega = \int_{\tilde{X}/C'} \omega \wedge \bar{\alpha} = \int_{\tilde{X}/C'} \omega \wedge *\omega = ||\omega||^2_{L^2(\tilde{X}/C')} \neq 0. \tag{5.18}$$

□

We can now prove the proposition 5.2. Let us briefly describe the idea. We consider the iterates F'^k of $F' : \tilde{X}/C' \to \tilde{X}/C'$. As F' is homotopic to the identity map, $F'^k(Z')$ is homologous to Z' and if ω is the harmonic form of the corollary 5.5 we have

$$\int_{Z'} (F'^k)^*(\omega) = \int_{Z'} \omega = a \neq 0. \tag{5.19}$$

We don't know if $F'^k(Z')$ converges or stays in a compact subset of $\tilde{X}/C'$ but we will show that $F'^k(Z')$ cannot entirely diverge in $\tilde{X}/C'$ and that there exists a $z' \in Z'$ such that $F'^k(z')$ subconverges to a point $x \in \tilde{X}/C'$ with $|Jac_{n-1}F'(x)| = 1$.

¿From (5.19) one gets

$$0 < |a| = |\int_{Z'} (F'^k)^*(\omega)| \leq \int_{Z'} |Jac_{TZ'} F'^k(z)|.||\omega(F'^k(z)||dz \tag{5.20}$$

where

$$|Jac_{TZ'} F'^k(z)| = ||DF'^k(z)(u_1) \wedge DF'^k(z)(u_2) \wedge ... \wedge DF'^k(z)(u_{n-1})||$$

and $(u_1, .., u_{n-1})$ is an orthonormal basis of $T_z(Z')$.

Let us define $\mathcal{B} = \{z \in Z', |JacF'^k_{TZ'}(z)|$ *does not converge to 0* $\}$.

For $z \in Z'$ we define the sequence z_k by $z_0 = z$ and $z_k = F'(z_{k-1}) = F'^k(z) \in \tilde{X}/C'$.

Lemma 5.6. *There exists $z \in \mathcal{B}$ and a subsequence z_{k_j} such that z_{k_j} converges to a point $x \in \tilde{X}/C'$ with $|Jac_{n-1}(x)| = 1$.*

Proof :

We first remark that $\lim_{x\to\infty} ||\omega(x)|| = 0$. This follows the following facts: ω is harmonic, $\omega \in L^2(\tilde{X}/C')$ and the injectivity radius of $\tilde{X}/C'$ is bounded below by a positive constant.

Let us assume that for all $z \in \mathcal{B}$ the sequence z_k diverges in $\tilde{X}/C'$. Then we have, for all $z \in \mathcal{B}$,

$$||\omega(z_k)|| = ||\omega(F'^k(z))|| \to 0 \tag{5.21}$$

whenever k tends to ∞ because of the previous remark.

On the other hand, as $||\omega(F'^k(z))|| \leq C$ and $|Jac_{n-1}F'^k| \leq 1$, it follows from (5.20)

$$\lim_{k\to\infty} |\int_{Z'} (F'^k)^*(\omega)| \leq$$

$$\lim_{k\to\infty} [\int_{\mathcal{B}} ||\omega(F'^k(z))||dz + C\int_{Z'-\mathcal{B}} |Jac_{Z'}F'^k(z)|dz = 0$$

which contradicts our assumption.

Thus there exists a point $z \in Z'$ such that

$$|Jac_{Z'}F'^k(z)| \to \alpha \neq 0 \tag{5.22}$$

and such that there exists a subsequence $z_{k_j} = F'^{k_j}(z)$ with

$$lim_{j\to\infty} z_{k_j} = x \in \tilde{X}/C'. \tag{5.23}$$

The property (5.22) comes from the fact that the sequence $|Jac_{Z'}F'^k(z)|$ doesn't tend to zero and is decreasing (because $|Jac_{n-1}F'| \leq 1$).

Let us define

$$E_0 = T_z Z', E_1 = DF'(z)(E_0)$$

and

$$E_k = DF'(z_{k-1})(E_{k-1}) \subset T_{z_k}(\tilde{X}/C').$$

As $z_{k_j} \to x$ we can assume, after extracting again a subsequence, that $E_{k_j} \to E \subset T_x(\tilde{X}/C')$. On the other hand we also have

$$|Jac_{Z'}F'^k(z)| = |Jac_{E_{k-1}}F'(z_{k-1})||Jac_{E_{k-2}}F'(z_{k-2})|...|Jac_{E_0}F'(z)| \tag{5.24}$$

We know that $|Jac_{E_k}F'(z_k)| = 1 - \epsilon_k$ where $0 \leq \epsilon_k < 1$. As $z \in \mathcal{B}$, we have

$$\lim_{k\to\infty} \pi_{j=1}^{k}(1 - \epsilon_j) = \alpha > 0$$

therefore $\lim_{k\to\infty} \epsilon_k = 0$ and by continuity we have $|Jac_E F'(x)| = 1$. $\square$

Now we can finish the proof of the step 1. Let consider a lift of x in $\tilde{X}$ and E in $T\tilde{X}$ that we again call x and E. Then we have $|Jac_E \tilde{F'}(x)| = 1$. $\square$

Let us remark that corollary 4.4 and (5.19) give another proof of the inequality $\delta \geq n-2$, which does not use the isosystolic inequality, ie. Theorem 3.3.

Step 2 : The weak tangent of $\partial\tilde{X}$ and Λ_C

We first recall the definition of the Gromov-distance on $\partial\tilde{X}$.

For two arbitrary points θ and θ' in $\partial\tilde{X}$ let us define

$$l(\theta, \theta') = inf\{t > 0 / dist(\alpha_\theta(t), \alpha_{\theta'}(t)) = 1\} \tag{5.25}$$

and

$$d(\theta, \theta') = e^{-l(\theta,\theta')} \tag{5.26}$$

then d is a distance on $\partial\tilde{X}$.

We now recall a few definitions following [4]. A complete metric space $(S, \bar{d})$ is a weak tangent of a metric space (Z, d) if there exist a point $0 \in S$, a sequence of points $z_k \in Z$ and a sequence of positive real numbers $\lambda_k \to \infty$ such that the sequence of pointed metric spaces $(Z, \lambda_k d, z_k)$ converges in the pointed Gromov-Hausdorff topology to $(S, \bar{d}, 0)$ where $(Z, \lambda_k d)$ stands for the set Z endowed with the rescaled metric $\lambda_k d$.

Let us recall that the sequence of metric spaces (Z_k, d_k, z_k) converges to $(S, \bar{d}, 0)$ in the pointed Gromov-Hausdorff topology if the following conditions hold, (cf. (B-B-I), definition 8.1.1).

Definition 5.7. *We say that the sequence of metric spaces (Z_k, d_k, z_k) converges to $(S, \bar{d}, o)$ if for any $R > 0$, $\epsilon > 0$ there exists k_0 such that for any $k \geq k_0$ there exists a (non necessary continuous) map $f : B(z_k, R) \to S$ such that*
(i) $f(z_k) = 0$,
for any two points x and y in $B(z_k, R)$,
(ii) $|\bar{d}(f(x), f(y)) - d_k(x, y)| \leq \epsilon$,
and
(iii) the ϵ-neighborhood of the set $f(B(z_k, R)$ contains $B(0, R - \epsilon)$.

In the previous definition, $B(z_k, R)$ stands for the ball of radius R centered at the point z_k in (Z_k, d_k).

For a metric space (Z, d) we will denote $WT(Z, d)$ the set of weak tangents of (Z, d).

Let Γ a cocompact group of isometries of $(\tilde{X}, \tilde{g})$ a n-dimensional Cartan Hadamard manifold of sectional curvature $K_{\tilde{g}} \leq -1$ and C' a subgroup of Γ. The limit set Λ_Γ of Γ is the full boundary $\partial\tilde{X}$ namely a topological $(n-1)-$dimensional sphere S^{n-1}. We endow $\partial\tilde{X}$ with the Gromov distance d defined in (5.26). In [4] Lemma 5.2, M.Bonk and B.Kleiner show among other properties the following

Lemma 5.8. *For any weak tangent space* $(S, \bar{d})$ *in* $WT((\partial\tilde{X}, d))$*,* S *is homeomorphic to* $\partial\tilde{X}$ *less a point, thus to* $\mathbb{R}^{n-1}$*.*

In fact the crucial asumption in the above lemma, coming from the cocompacness of Γ, is the property that any triple of points in $\partial\tilde{X}$ can be uniformly separated by an element of Γ, ie. there is $\delta > 0$ such that for any three points $\theta_1, \theta_2, \theta_3 \in \partial\tilde{X}$ there exists a $\gamma \in \Gamma$ such that $d(\gamma\theta_i, \gamma\theta_j) \geq \delta$ for all $1 \leq i \neq j \leq 3$. Following the argument of M.Bonk and B.Kleiner one can show that if C' is a subgroup of Γ such that one weak tangent $(S, \bar{d})$ of $(\Lambda_{C'}, d)$ is in $WT(\partial\tilde{X}, d)$ and enough triples of points of $\Lambda_{C'}$ can be uniformly separated by elements of C', then S is homeomorphic to $\Lambda_{C'}$ less a point. In particular, $\Lambda_{C'}$ is homeomorphic to $\partial\tilde{X}$.

Lemma 5.9. *Let* $\mathcal{L} \subset \partial\tilde{X}$ *be a closed* C'*-invariant set and* $\theta_0 \in \mathcal{L}$*. We assume that there exist a sequence of positive real numbers* $\lambda_k \to \infty$ *such that the sequence of pointed metric spaces* $(\mathcal{L}, \lambda_k d, \theta_0)$ *converges in the pointed Gromov-Hausdorff topology to* $(S, \bar{d}, 0)$ *where* $(S, \bar{d}, 0)$ *is a weak tangent of* $(\partial\tilde{X}, d)$*. We also assume that there exist positive constants* C *and* δ*, a sequence of points* $\theta_0^k = \theta_0, \theta_1^k, \theta_2^k \in \mathcal{L}$ *and a sequence of elements* $\gamma_k \in C'$ *such that* $C^{-1}\lambda_k^{-1} \leq d(\theta_i^k, \theta_j^k) \leq C\lambda_k^{-1}$ *and* $d(\gamma_k\theta_i^k, \gamma_k\theta_j^k) \geq \delta$ *for all* $0 \leq i \neq j \leq 2$*. Then,* S *is homeomorphic to* $\mathcal{L}$ *less a point. In particular* $\mathcal{L}$ *is homeomorphic to* $\partial\tilde{X}$*.*

The proof of this lemma is postponed in the Appendix.

Step 3 : The limit set $\Lambda_{C'}$ of C' and the limit set Λ_C of C are equal to a topological equator.

We have shown in step 1 that Λ_C is a subset of some topological equator $E(\infty)$.

Let $o \in \tilde{X}$ and $E \subset T_o\tilde{X}$ be such that $|Jac_E\tilde{F}(o)| = 1$ and $E(\infty)$ is the equator associated to E.

Recall that there exists a subgroup C' of C which globally preserves an hypersurface $\tilde{Z}' \subset \tilde{X}$ and that $\tilde{Z}'/C' \subset \tilde{X}/C'$ is compact. Furthermore $\tilde{Z}'$ separates $\tilde{X}$ into two connected components $\tilde{U}$ et $\tilde{V}$. We can assume that $\tilde{U}$ and $\tilde{V}$ are globally invariant by C' after having replaced C' by an index 2 subgroup.

The limit set $\Lambda_{C'}$ of C' is contained in Λ_C, therefore $\Lambda_{C'} \subset E(\infty)$.

We will show that $\Lambda_{C'} = E(\infty)$.

For any subset $W \in \tilde{X}$ we define the boundary at infinity ∂W of W by

$$\partial W = Cl(W) \cap \partial \tilde{X} \tag{5.27}$$

where $Cl(W)$ stands for the closure of W in $\tilde{X} \cup \partial \tilde{X}$.

As $\tilde{Z}'/C'$ is compact, $\tilde{Z}'$ is at bounded distance of the orbit $C'z$ of some point z in $\tilde{Z}'$, thus

$$Cl(\tilde{Z}') \cap \partial \tilde{X} = \Lambda_{C'} \tag{5.28}$$

By definition we have $\Lambda_{C'} \subset \partial \tilde{U}$ and $\Lambda_{C'} \subset \partial \tilde{V}$.

Lemma 5.10. *Let us assume that $\Lambda_{C'} \neq E(\infty)$, then either $\partial \tilde{U} = \Lambda_{C'}$ or $\partial \tilde{V} = \Lambda_{C'}$.*

Proof: We know that $\Lambda_{C'} \subset \partial \tilde{U}$ and $\Lambda_{C'} \subset \partial \tilde{V}$.

Let us assume that the conclusion of the lemma is not true, so there is $\zeta \in \partial \tilde{U} - \Lambda_{C'}$ and $\theta \in \partial \tilde{V} - \Lambda_{C'}$.

As $\Lambda_{C'} \subset E(\infty)$, $\Lambda_{C'} \neq E(\infty)$ and $\partial \tilde{X}$ is a sphere, any two points of $\partial \tilde{X} - \Lambda_{C'}$ can be joined by a continuous path contained in $\partial \tilde{X} - \Lambda_{C'}$ and so does ζ and θ, joined by such a path α.

The set $Z' \cup \Lambda_{C'}$ is a closed subset of $\tilde{X} \cup \partial \tilde{X}$ thus there is an open connected neighborhood W of α in $\tilde{X} \cup \partial \tilde{X}$ contained in the complementary of $Z' \cup \Lambda_{C'}$.

As ζ and θ can be approximated by points in $\tilde{U}$ and $\tilde{V}$ respectively there exist points $x \in \tilde{U} \cap W$ and $y \in \tilde{V} \cap W$ that can be joined by a continuous path by connectedness of W, which leads to a contradiction. □

Remark 5.11. *In fact, we are going to show that under the assumption $\delta(C') = n - 2$, it is impossible to have $\partial \tilde{U} = \Lambda_{C'}$ or $\partial \tilde{V} = \Lambda_{C'}$.*

For any $x \in \tilde{X}$ and $\theta \in \partial \tilde{X}$ let us denote $HB(x, \theta)$ the open horoball centered at θ and passing through x.

Lemma 5.12. *Let us assume that $\partial \tilde{U} = \Lambda_{C'}$. Then there exist $\theta_0 \in \Lambda_{C'}$ and $z' \in \tilde{Z}'$ such that $HB(z', \theta_0) \subset \tilde{U}$.*

Proof : Let us recall that $\tilde{X}/C' - Z' = U \cup V$ where $U = \pi(\tilde{U})$ $V = \pi(\tilde{V})$ and $\pi : \tilde{X} \to \tilde{X}/C'$ is the projection .

We know that U and V are unbounded. Let x_n a sequence of points in U such that $dist(x_n, Z') \to \infty$. Let $z_n \in Z'$ such that $dist(x_n, Z') = dist(x_n, z_n)$. We consider a fundamental domain $D \subset \tilde{Z}'$ of C'. There exist lifts $\tilde{z}_n \in D$ and $\tilde{x}_n \in \tilde{U}$ such that $dist(\tilde{x}_n, Z') = dist(\tilde{x}_n, \tilde{z}_n)$ tends to infinity.

By compactness we can assume that a subsequence $\tilde{x}_{n_j}$ converges to a point $\theta_0 \in \partial \tilde{X}$ and $\tilde{z}_{n_j}$ also converges to a point $\tilde{z} \in \bar{D}$. Furthermore the sequence of open balls $B(\tilde{x}_{n_j}, dist(\tilde{x}_{n_j}, \tilde{z}_{n_j})) \subset \tilde{U}$ converges to the open horoball $HB(\theta_0, \tilde{z}) \in \tilde{U}$. □

Proposition 5.13. *$\Lambda_{C'} = E(\infty)$ and $\Lambda_C = E(\infty)$.*

We first describe the idea of the proof and next state some facts we will need in order to do it.

As $\Lambda_{C'} \subset \Lambda_C \subset E(\infty)$ the proposition boils down to proving that $\Lambda_{C'} = E(\infty)$. Let us assume $\Lambda_{C'} \neq E(\infty)$ and find a contradiction.

We will show that for any sequence θ_i converging to θ_0 the geodesic starting at a point o such that $|Jac_{n-1}\tilde{F}(o)| = 1$ and ending at θ_i crosses the hypersurface $\tilde{Z}'$ in a point z_i. For an **appropriate choice** of such a sequence θ_i (roughly speaking, the sequence θ_i is chosen to be converging to θ_0 "transversally to $\Lambda_{C'}$"), the shadow (defined below) projected from o through some geodesic ball $B(z_i, r)$ will not intersect $\Lambda_{C'}$. On the other hand this shadow has to meet the limit set $\Lambda_{C'}$ because of the shadow lemma of D. Sullivan, which leads to a contradiction.

Precisely, by Lemma (5.10) and Lemma (5.12) we know that $\partial\tilde{U} = \Lambda_{C'}$ and that there exists an open horoball $HB(\theta_0, \tilde{z}) \subset \tilde{U}$ centered at a point $\theta \in \Lambda_{C'}$, and whose closure contains a point $\tilde{z} \in \tilde{Z}'$.

Let $o \in \tilde{X}$ and $E \in T_o\tilde{X}$ an hyperplane such that $\tilde{F}(o) = o$, $|Jac_E\tilde{F}(o)| = 1$, and $\Lambda_{C'} \subset E(\infty)$ where $E(\infty)$ is the topological equator associated to E.

For each $\theta \in \partial\tilde{X}$ we denote by α_θ the geodesic starting from o and such that $\alpha_\theta(+\infty) = \theta$.

Let $\theta_i \in \partial\tilde{X} - E(\infty) = \partial\tilde{V} - \partial\tilde{U}$ be a sequence converging to θ_0. By continuity, for each i large enough, the geodesic α_{θ_i} spends some time inside the horoball $HB(\theta_0, \tilde{z}) \subset \tilde{U}$ and ends up inside $\tilde{V}$ because θ_i converges to θ_0 and θ_i belongs to $\partial\tilde{V} - \partial\tilde{U}$.

Thus α_{θ_i} eventually crosses $\tilde{Z}'$. Let $z_i \in \alpha_{\theta_i} \cap \tilde{Z}'$. As $\tilde{Z}'/C'$ is compact, there is an element $\gamma_i \in C'$ such that $z_i = \gamma_i(x_i)$ where x_i is a point in the closure $\bar{D}$ of a fundamental domain D for the action of C' on $\tilde{Z}'$. The points $\gamma_i(x_i)$ and $\gamma_i(o)$ stay at bounded distance because $dist(\gamma_i(x_i), \gamma_i(o)) = dist(x_i, o) \leq dist(o, D) + diamD$. In particular, $lim_{i\to\infty}\gamma_i(o) = \theta_0$.

We have proved the

Lemma 5.14. *Let $\theta_i \in \partial\tilde{X} - E(\infty)$ be a sequence which converges to θ_0. There exists a constant A such that for i large enough there exists $z_i \in \tilde{Z}' \cap \alpha_{\theta_i}$ and $\gamma_i \in C'$ such that $dist(z_i, \gamma_i(o)) \leq A$ and both z_i and $\gamma_i(o)$ converge to θ_0.*

Let x and y two points in $\tilde{X}$.

We define the shadow $\mathcal{O}(x, y, R) \subset \partial\tilde{X}$ of the ball $B(y, R)$ enlighted from the point x by

$$\mathcal{O}(x, y, R) = \{\alpha(+\infty)\} \tag{5.29}$$

where α runs through the set of geodesic rays starting from x and meeting $B(y, R)$.

Let $\{\mu_x\}_x$ be a family of Patterson measures associated to the discrete group C' with critical exponent $\delta' = \delta(C')$.

The following shadow lemma is due to D.Sullivan.

Lemma 5.15. [18], [13], [20]. *There exist positive constants C and R such that for any y in $\tilde{X}$, $\nu_y(\mathcal{O}(y, \gamma(y), R)) \geq Ce^{\delta' d(y,\gamma(y))}$*

Corollary 5.16. *Let z_i be defined in lemma (5.14), then we have $\mathcal{O}(o, z_i, R + A) \cap \Lambda_{C'} \neq \emptyset$ for i large enough.*

We now prove that for a good choice of θ_i, the shadow $\mathcal{O}(o, z_i, R + A)$ (with z_i associated to θ_i as in lemma 5.14) never meet $\Lambda_{C'}$ for all large i's, ie. for any $\theta \in \Lambda_{C'}$ the geodesic α_θ does not cross $B(z_i, R + A)$. We have no control on the radius R coming from the shadow lemma nor on the constant A but we will show

Proposition 5.17. *There exists a sequence $\theta_i \in \partial\tilde{X} - \Lambda_{C'}$ such that θ_i converges to θ_0 and*

$$lim_{i\to\infty} inf_{\theta\in\Lambda_{C'}} dist(z_i, \alpha_\theta) = +\infty$$

where $z_i = \tilde{Z}' \cap \alpha_{\theta_i}$ has been constructed in lemma (5.14).

Corollary 5.18. *For i large enough, $\mathcal{O}(o, z_i, R + A) \cap \Lambda_{C'} = \emptyset$.*

The corollary (5.16) and the corollary (5.18) lead to a contradiction, which ends the proof of the proposition (5.13).

The end of the paragraph is devoted to proving the proposition (5.17).

Lemma 5.19. *Let θ_i be a sequence of points in $\partial\tilde{X}$ converging to θ_0 and z_i constructed in lemma (5.14). Assume that $liminf_{i\to\infty} inf_{\theta\in\Lambda_{C'}} dist(z_i, \alpha_\theta) = C < +\infty$ then $lim_{i\to\infty} \frac{d(\theta_i, \Lambda_{C'})}{d(\theta_i, \theta_0)} = 0$.*

Proof : We first show that

$$\lim_{i\to\infty} dist(z_i, \alpha_{\theta_0}) = \infty \tag{5.30}$$

Recall that, for any $z \in \tilde{X}$ and $\theta \in \partial\tilde{X}$, $B(z, \theta)$ equals the decreasing limit as t tends to infinity of $dist(z, \alpha_\theta(t)) - dist(o, \alpha_\theta(t))$ where $\alpha_\theta(t)$ is the geodesic ray joigning o to θ. Therefore, as the points $z_i \in \tilde{Z}$ belongs to the complementary of the fixed horoball $HB(\tilde{z}, \theta_0)$, we have,

$$dist(z_i, \alpha_{\theta_0}(T_i) \geq T_i + B(\tilde{z}, \theta_0) \tag{5.31}$$

where $dist(z_i, \alpha_{\theta_0}(T_i)) = dist(z_i, \alpha_{\theta_0})$.

On the other hand, as z_i tends to θ_0, T_i tends to infinity so (5.30) is proven.

Let t_i be such that $z_i = \alpha_{\theta_i}(t_i)$. By (5.30) we have

$$\lim_{i\to\infty} dist(\alpha_{\theta_i}(t_i), \alpha_{\theta_0}(t_i)) = \infty. \tag{5.32}$$

Let u_i be such that

$$\mathrm{dist}(\alpha_{\theta_\mathrm{i}}(\mathrm{u_i}), \alpha_{\theta_0}(\mathrm{u_i})) = 1, \tag{5.33}$$

then in particular $u_i \leq t_i$ for i large enough and by the triangle inequality we have

$$dist(\alpha_{\theta_i}(t_i), \alpha_{\theta_0}(t_i)) \leq 2(t_i - u_i) + 1. \tag{5.34}$$

By (5.32) we get

$$\lim_{i\to\infty} (t_i - u_i) = +\infty. \tag{5.35}$$

Let us assume there exists a sequence $\theta'_i \in \Lambda_{C'}$ and a constant C such that

$$dist(z_i, \alpha_{\theta'_i}) \leq C < +\infty. \tag{5.36}$$

We can assume that $C \geq 1$.
Let v_i be such that

$$dist(z_i, \alpha_{\theta'_i}) = dist(z_i, \alpha_{\theta'_i}(v_i)). \tag{5.37}$$

By triangle inequality,

$$|t_i - v_i| \leq C \tag{5.38}$$

and,

$$dist(\alpha_{\theta'_i}(t_i), \alpha_{\theta_i}(t_i)) \leq 2C. \tag{5.39}$$

On the other hand, as the curvature of $\tilde{X}$ is bounded above by -1, a classical comparison theorem gives for any $t \in [0, t_i]$,

$$sinh\left(\frac{dist(\alpha_{\theta'_i}(t), \alpha_{\theta_i}(t))}{2}\right) \leq sinhC.\frac{sinht}{sinht_i}. \tag{5.40}$$

Let s_i be such that

$$dist(\alpha_{\theta'_i}(s_i), \alpha_{\theta_i}(s_i)) = 1. \tag{5.41}$$

There are two cases. Either $s_i \geq t_i$ or $s_i < t_i$. If $s_i < t_i$, we get from (5.39) and (5.40) the existence of a constant A such that for any i,

$$s_i \geq t_i - A, \tag{5.42}$$

and this inequality also holds when $s_i \geq t_i$.

¿From (5.42) we get

$$\frac{d(\theta_i, \theta'_i)}{d(\theta_i, \theta_0)} = e^{-s_i+u_i} \leq e^A e^{-t_i+u_i}, \tag{5.43}$$

therefore, thanks to (5.35) we obtain

$$lim_{i\to\infty} \frac{d(\theta_i, \theta'_i)}{d(\theta_i, \theta_0)} = 0. \tag{5.44}$$

which ends the proof of lemma (5.19). □

Lemma 5.20. *Let us assume that for every sequence θ_i of points in $\partial\tilde{X}$ converging to θ_0, $lim_{i\to\infty}\frac{d(\theta_i,\Lambda_{C'})}{d(\theta_i,\theta_0)} = 0$. Let $\lambda_k \to \infty$ be such that the sequence of spaces $(\partial\tilde{X}, \lambda_k d, \theta_0)$ converges to a space $(S, \bar{d}, 0)$ in the pointed Gromov-Hausdorff topology, then the sequence of spaces $(\Lambda_{C'}, \lambda_k d, \theta_0)$ also converges to $(S, \bar{d}, 0)$.*

Proof : Let us define

$$r(\epsilon) =: sup\{\frac{d(\theta, \Lambda_{C'})}{d(\theta, \theta_0)}, \theta \neq \theta_0, d(\theta, \theta_0) \leq \epsilon\}.$$

The assumption says that

$$lim_{\epsilon\to 0} r(\epsilon) = 0. \tag{5.45}$$

For an arbitrary metric space (Y, d) and Y' a subset of Y, let us denote $B_{(Y,d)}(y, R)$ the closed ball of (Y, d) of radius R centered at $y \in Y$, and $\mathcal{U}_\epsilon^{(Y,d)}(Y')$ the ϵ-neighborhood of Y' in (Y, d). For a metric space (Y, d) and a positive number λ, let us denote λY the rescaled space $(Y, \lambda d)$.

By definition of the function r, we have for any R,

$$B_{\lambda_k\partial\tilde{X}}(\theta_0, R) \subset \mathcal{U}_{\epsilon_k}^{\lambda_k\partial\tilde{X}} B_{\lambda_k\Lambda_{C'}}(\theta_0, R + \epsilon_k)$$

$$\subset B_{\lambda_k\partial\tilde{X}}(\theta_0, R + 2\epsilon_k). \tag{5.46}$$

where $\epsilon_k =: Rr(R/\lambda_k)$.

Let us fix $\alpha > 0$. By definition 5.7, for any $R > 0$, $\epsilon > 0$, there exist a map $f : B_{\lambda_k\partial\tilde{X}}(\theta_0, R + \alpha) \to S$ such that for $k \geq k_0$,
(i) $f(\theta_0) = 0$,
for any two points x and y in $B_{\lambda_k\partial\tilde{X}}(\theta_0, R + \alpha)$,

(ii) $|\bar{d}(f(x), f(y)) - \lambda_k d(x, y)| \le \epsilon$,
and
(iii) $B_{(S,\bar{d})}(0, R+\alpha-\epsilon) \subset \mathcal{U}_\epsilon^{(S,\bar{d})} f(B_{\lambda_k \partial\tilde{X}}(\theta_0, R+\alpha))$.

Moreover let us prove:
(iv) $B_{(S,\bar{d})}(0, R-2\epsilon) \subset \mathcal{U}_\epsilon^{(S,\delta)} f(B_{\lambda_k \partial\tilde{X}}(\theta_0, R))$.
Indeed, let $z \in B_{(S,\bar{d})}(0, R-2\epsilon)$. By (iii), there exists $\theta \in B_{\lambda_k \partial\tilde{X}}(\theta_0, R+\alpha)$ such that $\bar{d}(z, f(\theta)) \le \epsilon$. Since $\bar{d}(z, 0) \le R-2\epsilon$, we thus deduce from triangle inequality $\bar{d}(f(\theta), 0) \le R-\epsilon$, and therefore we get, thanks to (i) and (ii), $\lambda_k d(\theta, \theta_0) \le R$. □

By (5.45), for ϵ small enough, there exists $k_1 \ge k_0$ such that for any $k \ge k_1$, then $2\epsilon_k \le \epsilon$ and

$$B_{\lambda_k \partial\tilde{X}}(\theta_0, R+2\epsilon_k) \subset B_{\lambda_k \partial\tilde{X}}(\theta_0, R+\alpha).$$

Therefore, by (5.46) and the above properties (i),(ii),(iii) and (iv) of the map f, and the triangle inequality we get,

$$B_{(S,\bar{d})}(0, R-2\epsilon) \subset \mathcal{U}_\epsilon^{(S,\bar{d})} f(B_{\lambda_k \partial\tilde{X}}(\theta_0, R))$$

$$\subset \mathcal{U}_\epsilon^{(S,\bar{d})} f(\mathcal{U}_{\epsilon_k}^{\lambda_k \partial\tilde{X}} B_{\lambda_k \Lambda_{C'}}(\theta_0, R+\epsilon_k))$$

$$\subset \mathcal{U}_{2\epsilon+\epsilon_k}^{(S,\bar{d})} f(B_{\lambda_k \Lambda_{C'}}(\theta_0, R+\epsilon_k)).$$

About the second inclusion above let us remark that the set $\mathcal{U}_{\epsilon_k}^{\lambda_k \partial\tilde{X}} B_{\lambda_k \Lambda_{C'}}(\theta_0, R+\epsilon_k)$ is contained in $B_{\lambda_k \partial\tilde{X}}(\theta_0, R+2\epsilon_k) \subset B_{\lambda \partial\tilde{X}}(\theta_0, R+\alpha)$, so that we can apply f to this set.

¿From the above inclusions we obtain

$$B_{(S,\bar{d})}(0, R-3\epsilon) \subset \mathcal{U}_{2\epsilon+\epsilon_k}^{(S,\bar{d})} f(B_{\lambda_k \Lambda_{C'}}(\theta_0, R))$$

which implies the convergence of $(\Lambda_{C'}, \lambda_k d, \theta_0)$ to $(S, \bar{d}, 0)$.
□

Corollary 5.21. *Let us assume that for every sequence θ_i of points in $\partial\tilde{X}$ converging to θ_0 and z_i the sequence of points constructed in lemma (5.14), $liminf_{i\to\infty} inf_{\theta\in\Lambda_{C'}} dist(z_i, \alpha_\theta) < +\infty$. Let $\lambda_k \to \infty$ be such that the sequence of spaces $(\partial\tilde{X}, \lambda_k d, \theta_0)$ converges to the space $(S, \delta, 0)$ in the pointed Gromov-Hausdorff topology, then the sequence of spaces $(\Lambda_{C'}, \lambda_k d, \theta_0)$ also converges to $(S, \delta, 0)$.*

We will show now that there exist a sequence of points $\theta_1^k, \theta_2^k \in \Lambda_{C'}$ converging to θ_0, such that the mutual distances $d(\theta_1^k, \theta_2^k)$, $d(\theta_1^k, \theta_0)$, $d(\theta_2^k, \theta_0)$ is tending to zero at the same rate, and the triple $\theta_1^k, \theta_2^k, \theta_0$ can be uniformly separated by elements $\gamma_k \in C'$.

Lemma 5.22. *Assume that every weak tangent of* $(\partial\tilde{X}, d)$ *at* θ_0 *belongs to* $WT(\Lambda_{C'}, d)$*, then there exist positive constants* c, δ*, a sequence* ϵ_k *tending to* 0 *when* k *tends to* ∞*, a sequence* $\gamma_k \in C'$*, a sequence of points* $\theta_1^k, \theta_2^k \in \Lambda_{C'}$ *such that for* $i = 1, 2$*,*
$c^{-1}\epsilon_k \leq d(\theta_1^k, \theta_2^k) \leq c\epsilon_k$*,*
$c^{-1}\epsilon_k \leq d(\theta_i^k, \theta_0)) \leq c\epsilon_k$ *and*
$d(\gamma_k\theta_1^k, \gamma_k\theta_2^k) \geq \delta$*,* $d(\gamma_k\theta_i^k, \gamma_k\theta_0)) \geq \delta$*.*

Proof : For any $x \in \tilde{X} \cup \partial\tilde{X}$ and $y \in \tilde{X} \cup \partial\tilde{X}$ let us define $\alpha_{x,y}$ the geodesic ray joining x and y. Let $o \in \tilde{X}$ and $E \in T_o\tilde{X}$ be such that $|Jac_E\tilde{F}(o)| = 1$ and $E(\infty)$ the equator associated to E. Let $\gamma_k \in C'$ be a sequence such that $\gamma_k(o)$ converges to the point θ_0 where $\theta_0 \in \Lambda_{C'}$ is the point coming from lemma 5.12. In particular, according to that lemma, there exist a point $\tilde{z} \in \tilde{Z}'$ such that the hypersurface $\tilde{Z}'$ is contained in the complementary of the open horoball $HB(\tilde{z}, \theta_0)$. We define $D := dist(\tilde{z}, o)$. As $\tilde{Z}'$ lies outside the open horoball $HB(\tilde{z}, \theta_0)$, the points $\gamma_k(o)$ belong to the complementary of the open horoball $HB(\alpha_{\tilde{z},\theta_0}(D), \theta_0)$. By standard triangle comparison argument (comparison with the hyperbolic case) the angle $Angle(\alpha_{\gamma_k(o),\theta_0}, \alpha_{\gamma_k(o),o})$ between the two geodesic rays $\alpha_{\gamma_k(o),\theta_0}$ and $\alpha_{\gamma_k(o),o}$ satisfies :

$$lim_{k\to\infty}Angle(\alpha_{\gamma_k(o),\theta_0}, \alpha_{\gamma_k(o),o}) = 0. \tag{5.47}$$

By equivariance we have $\Lambda_{C'} \subset (\gamma_k E)(\infty)$ where $\gamma_k E \subset T_{\gamma_k(o)}\tilde{X}$. For any $v \in T\tilde{X}$ let α_v be the geodesic ray such that $\dot{\alpha}_v(0) = v$. Let us denote by u_k the unit vector in $\gamma_k E$ such that $\alpha_{u_k}(+\infty) = \theta_0$ and let us choose some $w_k \in \gamma_k E$ such that $< u_k, w_k >= 0$ (this is possible because $n - 1 \geq 2$.

We claim now that there exist $v_k \in \gamma_k E$ such that the angle between v_k and w_k is not too far from 0 or π, namely

$$| < v_k, w_k > | \geq \frac{1}{(n-1)^{1/2}}, \tag{5.48}$$

and $\alpha_{v_k}(+\infty) \in \Lambda_{C'}$ or $\alpha_{v_k}(-\infty) \in \Lambda_{C'}$.

Let us prove this claim.

According to Proposition 5.1 and to (5.11), the restriction to $\gamma_k E$ of the quadratic form $h(u) = \int DB(\gamma_k(o), \theta)(u)^2 d\mu_{\gamma_k(o)}(\theta)$ verifies

$$h_{\gamma_k E}(u) = \frac{||u||^2}{n-1}. \tag{5.49}$$

Therefore, if for all $u \in \gamma_k(E)$ such that $\alpha_u(+\infty) = \theta \in \Lambda_{C'}$ we had $| < u, w_k > | < \frac{1}{(n-1)^{1/2}}$, then one would get $h(w_k) < \frac{1}{n-1}$, which contradicts (5.49) and proves the claim.

In particular the angle between u_k and v_k is not too far from $\pi/2$ for k large enough, ie.

$$|< u_k, v_k >| \leq \Big(\frac{n-2}{n-1}\Big)^{1/2}, \tag{5.50}$$

and thanks to (5.47), we have for k large enough

$$|< \dot{\alpha}_{\gamma_k(0),o}(0), v_k >| \leq \Big(\frac{n-\frac{3}{2}}{n-1}\Big)^{1/2}. \tag{5.51}$$

Let us now assume for example that $\theta_k = \alpha_{v_k}(+\infty) \in \Lambda_{C'}$. Let us show that

$$lim_{k\to\infty} d(\theta_0, \theta_k) = 0. \tag{5.52}$$

Assume that (5.52) is not true. Then, one can assume after extracting a subsequence that θ_k converges to $\theta \neq \theta_0$. Therefore the geodesic rays $\alpha_{\gamma_k(0),o}$ and α_{v_k} would converge to the geodesics $\alpha_{\theta_0,o}$ and $\alpha_{\theta_0,\theta}$ and thus the angle $Angle(\alpha_{\gamma_k(0),o}, \alpha_{v_k})$ would converge to 0. But this would contradict (5.51).

Let us now denote $\epsilon_k =: d(\theta_k, \theta_0)$. According to (5.52), $lim_{k\to\infty}\epsilon_k = 0$. We now consider the following sequence of pointed metric space $(\partial\tilde{X}, \epsilon_k^{-1} d, \theta_0)$, a subsequence of which being converging to some metric space (S, δ), cf[]. For convenience we still denote by the same index k the subsequence. By the corollary 5.21, the sequence $(\Lambda_{C'}, \epsilon_k^{-1} d, \theta_0)$ also converges to (S, δ). According to lemma 5.8, the space S is homeomorphic to $\mathbb{R}^{n-1}$. In particular there exist a sequence of points $\theta'_k \in \Lambda_{C'}$ and a constant c such that

$$c^{-1}\epsilon_k \leq d(\theta_k, \theta'_k) \leq c\epsilon_k, \tag{5.53}$$

$$c^{-1}\epsilon_k \leq d(\theta'_k, \theta_0)) \leq c\epsilon_k. \tag{5.54}$$

Thus, the points $\theta_1^k = \theta_k$ and $\theta_2^k = \theta'_k$ satisfy the two first properties of lemma 5.22.

In order to complete the proof of lemma 5.22, we will show that the elements $\eta_k =: \gamma_k^{-1}$ uniformly separate θ_0, θ_1^k and θ_2^k.

Thanks to (5.50) the angle at $\gamma_k(o)$ between θ_1^k and θ_0 is uniformly bounded away from 0 and π and so does the angle at o between $\gamma_k^{-1}(\theta_1^k)$ and $\gamma_k^{-1}(\theta_0)$. Therefore, as the angle is Hölder-equivalent to the distance d, cf. [10], there is a constant c such that

$$d(\gamma_k^{-1}(\theta_1^k), \gamma_k^{-1}(\theta_0)) \geq c. \tag{5.55}$$

Now the cocompact group Γ acts uniformly quasi-conformally on $(\partial\tilde{X}, d)$, ([4] and [19] Theorem 5.2), and so does $C' \subset \Gamma$, therefore

$$d(\gamma_k^{-1}(\theta_1^k), \gamma_k^{-1}(\theta_2^k)) \geq c, \tag{5.56}$$

and

$$d(\gamma_k^{-1}(\theta_2^k), \gamma_k^{-1}(\theta_0)) \geq c. \tag{5.57}$$

which ends the proof of lemma 5.22. □

Proof of Proposition 5.17 :

Let us assume that for every sequence θ_i of points in $\partial\tilde{X}$ converging to θ_0, $liminf_{i\to\infty} inf_{\theta\in\Lambda_{C'}} dist(z_i, \alpha_\theta) < +\infty$, then by corollary 5.21 and lemma 5.22 there exist a positive constant c, a sequence ϵ_k tending to 0 when k tends to ∞, a sequence $\gamma_k \in C'$, a sequence of points $\theta_1^k, \theta_2^k \in \Lambda_{C'}$ such that
$c^{-1}\epsilon_k \leq d(\theta_1^k, \theta_2^k) \leq c\epsilon_k$,
$c^{-1}\epsilon_k \leq d(\theta_i^k, \theta_0)) \leq c\epsilon_k$ and
$d(\gamma_k\theta_1^k, \gamma_k\theta_2^k) \geq \delta$, $d(\gamma_k\theta_i^k, \gamma_k\theta_0)) \geq \delta$.

Applying lemma 5.9 for $\mathcal{L} = \Lambda_{C'}$ and $\lambda_k = \epsilon_k^{-1}$ we conclude that $\Lambda_{C'}$ is homeomorphic to $\partial\tilde{X}$, which is impossible because $\Lambda_{C'}$ is contained in a topological equator $E(\infty)$. □

Step 4 : C' and C are convex cocompact.

We first define convex cocompactness. For a discrete group C of isometries acting on a Cartan Hadamard manifold of negative sectional curvature with limit set Λ_C, one defines the geodesic hull $\mathcal{G}(\Lambda_C)$ of Λ_C as the set of all geodesics both ends of whose belong to Λ_C.

The geodesic hull of Λ_C is a C invariant set. One says that C is convex cocompact if $\mathcal{G}(\Lambda_C)/C$ is compact.

Lemma 5.23. *C' is convex cocompact.*

Proof : Let us denote $\pi : \tilde{X} \to \tilde{X}/C'$ the projection. Assume that C' is not convex cocompact. Then, there exist a sequence $x_n \in \mathcal{G}(C')$ such that x_n tends to infinity. In particular $dist(x_n, Z') \to +\infty$, where $Z' = \tilde{Z}'/C'$ is the compact hypersurface which separates $\tilde{X}/C'$ in two unbounded connected components. There exist lifts $\tilde{x}_n$ of x_n such that

$$\tilde{x}_n \to \theta_0 \in \Lambda_{C'} \tag{5.58}$$

$$dist(\tilde{x}_n, \tilde{Z}') = dist(\tilde{x}_n, \tilde{z}_n) \tag{5.59}$$

where $\tilde{z}_n \in \tilde{Z}'$ is bounded. Therefore there exist $\tilde{z} \in \tilde{Z}'$
such that $HB(\tilde{z}, \theta_0) \subset \tilde{U}$, where $\tilde{U}$ is one of the two connected components of $\tilde{X} - \tilde{Z}'$, the other being $\tilde{V}$.

We recall that $\mathcal{M}$, $\mathcal{N}$ are the two connected components of $\partial\tilde{X} - \Lambda_{C'}$. We also have $\partial\tilde{Z}' = \Lambda_{C'} = E(\infty)$, and after possibly replacing C' by an index two subgroup, we can assume that C' preserves $\tilde{U}$ and $\tilde{V}$.

Claim : There are the two following cases.

Either one of the two boundaries $\partial\tilde{U}$ or $\partial\tilde{V}$ is equal to $\Lambda_{C'}$ (in this case the other boundary is equal to $\partial\tilde{X}$), or $\partial\tilde{U} = \bar{\mathcal{M}}$ and $\partial\tilde{V} = \bar{\mathcal{N}}$, where $\bar{\mathcal{M}}$ and $\bar{\mathcal{N}}$ are the closure of $\mathcal{M}$ and $\mathcal{N}$.

Let us prove the claim. We first remark that if there exist $\theta \in \partial\tilde{U} \cap \mathcal{M}$, then $\mathcal{M} \subset \partial\tilde{U}$. Namely, let ξ be any other point in $\mathcal{M}$ and α a continuous path in $\mathcal{M}$ joining θ and ξ. Since the set $\tilde{Z}' \cup \Lambda_{C'}$ is a closed subset in $\tilde{X} \cup \partial\tilde{X}$, there exist an open connected neighborhood W of α in $\tilde{X} \cup \partial\tilde{X}$ contained in the complementary of $\tilde{Z}' \cup \Lambda_{C'}$. Therefore, as $W \cap \tilde{U} \neq \emptyset$, we have $W \cap \tilde{X} \subset \tilde{U}$ and $\xi \in \partial\tilde{U}$. Let us assume that neither $\partial\tilde{U}$ nor $\partial\tilde{V}$ is equal to $\Lambda_{C'}$. Then, each boundary $\partial\tilde{U}$ and $\partial\tilde{V}$ contains $\mathcal{M}$ or $\mathcal{N}$. But on the other hand, since the set $\tilde{Z}' \cup \Lambda_{C'}$ is closed, $(\partial\tilde{U} - \Lambda_{C'}) \cap (\partial\tilde{V} - \Lambda_{C'}) = \emptyset$ thus we have $\partial\tilde{U} = \mathcal{M}$ and $\partial\tilde{V} = \mathcal{N}$ or the other way around and the claim is proved.

Case 1 : $\partial\tilde{U} = \Lambda_{C'}$ and $\partial\tilde{V} = \partial\tilde{X}$ or the other way around.

In this case, we are in the situation of the step 3, which leads to a contradiction, cf. remark 5.11.

Case 2 : $\partial\tilde{U} = \bar{\mathcal{M}}$ and $\partial\tilde{V} = \bar{\mathcal{N}}$.

In that case, assuming C' is not convex-cocompact, there exist an open horoball $HB(\theta_0, \tilde{z}) \subset \tilde{U}$ where $\theta_0 \in \Lambda_{C'}$, $\tilde{z} \in \tilde{Z}'$, $\partial\tilde{U} = \bar{\mathcal{M}}$ and $\partial\tilde{V} = \bar{\mathcal{N}}$. We will find a contradiction in a similar way as in case 1, ie. step 3. We consider a point $o \in \tilde{X}$ and an hyperplane $E \subset T_o\tilde{X}$ such that $|Jac_E\tilde{F}(o)| = 1$ and $\Lambda_{C'} = E(\infty)$.

Let $\theta_i \in \mathcal{N}$ be a sequence which converge to θ_0. By continuity, for i large enough, the geodesic ray α_{o,θ_i} spends some time in $HB(\theta_0, \tilde{z}) \subset \tilde{U}$ and ends up in $\tilde{V}$ because θ_i converges to θ_0 and θ_i belongs to $\mathcal{N} = \partial\tilde{V} - \Lambda_{C'}$. Therefore, α_{o,θ_i} eventually crosses $\tilde{Z}'$. Let z_i be some point in $\tilde{Z}' \cap \alpha_{o,\theta_i}$.

We will prove the following Proposition, similar to the Proposition 5.18,

Proposition 5.24. *There exist a sequence $\theta_i \in \mathcal{N}$ such that θ_i converges to θ_0 and*

$$lim_{i\to\infty} inf_{\theta\in\Lambda_{C'}} dist(z_i, \alpha_\theta) = +\infty$$

where $z_i \in \tilde{Z}' \cap \alpha_{\theta_i}$.

Remark 5.25. *The difference between the propositions 5.24 and 5.18 is that we are looking for a sequence $\theta_i \in \mathcal{N}$ instead of $\theta_i \in \partial\tilde{X} - \Lambda_{C'}$.*

Assuming the Proposition 5.24 we find a contradiction in the same way as in step 3. Namely, as $\tilde{Z}'/C'$ is compact, the points $z_i \in \tilde{Z}' \cap \alpha_{\theta_i}$ stay at bounded distance from the C'-orbit of a fixed point, say, o, thus there exist a constant $A > 0$ and elements $\gamma_i \in C'$ such that for any i,

$$dist(z_i, \gamma_i o) \leq A. \tag{5.60}$$

¿From (5.60) and the shadow lemma 5.15, we obtain $\mathcal{O}(o, z_i, R{+}A) \cap \Lambda_{C'} \neq \emptyset$, and on the other hand, from the proposition 5.24, we have $\mathcal{O}(o, z_i, R{+}A) \cap \Lambda_{C'} = \emptyset$, which gives the contradiction. It remains to prove the Proposition 5.24.

Proof of the proposition 5.24 : We argue by contradiction, like in the proof of the proposition 5.17. Let us assume that there exist a constant $C > 0$ such that for any sequence of points $\theta_i \in \mathcal{N}$ converging to θ_0, $lim_{i \to \infty} inf_{\theta \in \Lambda_{C'}} dist(z_i, \alpha_\theta) \leq C$, then by lemma 5.19, we have for any such sequence $\theta_i \in \mathcal{N}$

$$lim_{i \to \infty} \frac{d(\theta_i, \Lambda_{C'})}{d(\theta_i, \theta_0)} = 0. \tag{5.61}$$

The proof of the following lemma is the same as the proof of lemma 5.20.

Lemma 5.26. *Let us assume that for any sequence $\theta_i \in \mathcal{N}$ conveging to θ_0, $lim_{i \to \infty} \frac{d(\theta_i, \Lambda_{C'})}{d(\theta_i, \theta_0)} = 0$. Let $\{\lambda_k\}$ be a sequence of positive numbers tending to $+\infty$ such that the sequence of spaces $(\partial \tilde{X}, \lambda_k d, \theta_0)$ converges to a space $(S, \delta, 0)$ in the pointed Gromov-Hausdorff topology, then, $(\mathcal{M}, \lambda_k d, \theta_0)$ also converges to $(S, \delta, 0)$.*

Proof : Since $\Lambda_{C'} \subset \bar{\mathcal{M}}$, the assumption implies that $lim_{\epsilon \to 0} r(\epsilon) = 0$ where

$$r(\epsilon) = sup\{\frac{d(\theta, \mathcal{M})}{d(\theta, \theta_0)}, \theta \neq \theta_0, \theta \in \mathcal{N}, d(\theta, \theta_0) \leq \epsilon\}$$

and the proof goes the same way as in lemma 5.20 replacing $\Lambda_{C'}$ by $\mathcal{M}$. □

Similarly to the lemma 5.22, we have the

Lemma 5.27. *Let us assume that every weak tangent of $(\partial \tilde{X}, d)$ at θ_0 belongs to $WT((\mathcal{M}, d))$. There exist positive constant c, δ, a sequence ϵ_k tending to 0 when k tends to $+\infty$, a sequence of $\gamma_k \in C'$, a sequence of points $\theta_0^k = \theta_0, \theta_1^k, \theta_2^k \in \mathcal{M}$ such that for $i \neq j \in \{0, 1, 2\}$,*
$c^{-1}\epsilon_k \leq d(\theta_i^k, \theta_j^k) \leq c\epsilon_k$ and
$d(\gamma_k \theta_i^k, \gamma_k \theta_i^k) \geq \delta$.

We can now end the proof of the proposition 5.24. Let us assume that there exist a constant $C > 0$ such that for every sequence θ_i of points in $\mathcal{N}$ converging to θ_0, $lim_{i \to \infty} inf_{\theta \in \Lambda_{C'}} dist(\theta_i, \alpha_\theta) \leq C$, then by (5.61), lemma 5.26 and lemma 5.27, there exist a sequence ϵ_k tending to 0 when k tends to ∞, a sequence $\gamma_k \in C'$, a sequence of points $\theta_0^k = \theta_0, \theta_1^k, \theta_2^k \in \mathcal{M}$ such that for $i \neq j \in \{0, 1, 2\}$,

$c^{-1}\epsilon_k \leq d(\theta_i^k, \theta_j^k) \leq c\epsilon_k$ and
$d(\gamma_k\theta_i^k, \gamma_k\theta_i^k) \geq \delta$.
Applying the lemma 5.9 for $\mathcal{L} = \mathcal{M}$ and $\lambda_k = \epsilon_k^{-1}$, we conclude that $\mathcal{M}$ is homeomorphic to $\partial\tilde{X}$, which is impossible because $\partial\tilde{X}$ is a sphere, and $\mathcal{M}$ is homeomorphic to an hemisphere. This ends the proof of the proposition 5.24. □

Corollary 5.28. *C is convex cocompact.*

Proof : The subgroup C' of C is convex cocompact and the limit sets of C' and C coincide by step 3, therefore C is convex cocompact. □.

Step 5: C preserves a copy of the $(n-1)$-dimensional hyperbolic space $\mathbb{H}^{n-1}$ totally geodesically embedded in $\tilde{X}$.

From the steps 1-4, we know that the groups C and C' are convex cocompact, and that their limit set Λ_C and $\Lambda_{C'}$ are equal to a topological equator $E(\infty)$.

Let us consider the essential hypersurface $Z' \subset \tilde{X}/C'$. We will show that there exist a minimizing current representing the class of Z' in $H_{n-1}(\tilde{X}/C', \mathbb{R})$ and that this minimizing current lifts to a totally geodesic hypersurface embedded in $\tilde{X}$. We will then show that this totally geodesic hypersurface is eventually hyperbolic.

We work in $\tilde{X}/C'$ and consider the essential hypersurface $Z' \subset \tilde{X}/C'$. We will now prove that there exist a minimal current representing the class of Z' in $H_{n-1}(\tilde{X}/C', \mathbb{R})$. Let $\{Z_k\}$ be a minimizing sequence of currents homologous to Z'. The othogonal projection onto the convex core of $\tilde{X}/C'$ is distance nonincreasing and thus volume nonincreasing. Therefore we can assume that the Z_k's are in the the convex core of $\tilde{X}/C'$, which is compact. By [12] (5.5), the sequence $\{Z_k\}$ subconverges to a minimal current Z_∞ in $\tilde{X}/C'$. By [12] (8.2), Z_∞ is a manifold with possible singularities of codimension greater than or equal to 8. By corollary (4.4) and minimality we get that $|Jac_{n-1}\tilde{F}(x)| = 1$ at every regular points $x \in Z_\infty$. We will use the fact that $|Jac_{n-1}\tilde{F}(x)| = 1$ at every regular points $x \in Z_\infty$ in order to prove that Z_∞ is a totally geodesic hypersurface.

Lemma 5.29. *Let x and y two distinct points in $\tilde{X}$ and $E_x \subset T_x\tilde{X}$, $E_y \subset T_y\tilde{X}$ be such that $Jac_{n-1}\tilde{F}'(x) = Jac_{E_x}\tilde{F}'(x) = 1$ and $Jac_{n-1}\tilde{F}'(y) = Jac_{E_y}\tilde{F}'(y) = 1$. Then, the geodesic $\alpha_{x,y}$ (resp. $\alpha_{y,x}$) joining x and y (resp. y and x) satisfies $\dot{\alpha}_{x,y}(0) \in E_x$, (resp. $\dot{\alpha}_{y,x}(0) \in E_y$). In particular, $\alpha_{x,y}(+\infty)$ and $\alpha_{y,x}(+\infty)$ belong to $\Lambda_{C'}$.*

Proof: Let S_x and S_y be the unit spheres of E_x and E_y. For any unit tangent vector $u \in T_z\tilde{X}$ at some point z, we define $\theta_u \in \partial\tilde{X}$ by $\dot{\alpha}_{z,\theta_u}(0) = u$. By step 3, $\Lambda_{C'} = E_x(\infty) = E_y(\infty)$, therefore for every $u \in S_x$, $\theta_u \in \Lambda_{C'}$ and there exist $v \in E_y$ such that $\theta_u = \theta_v$. As E_y is a vector space, θ_{-v} belongs to

$\Lambda_{C'}$ therefore there exist $w \in E_x$ such that $\theta_w = \theta_{-v}$. The map $f : S_x \to S_x$ defined by $f(u) = w$ is a continuous map. The lemma then boils down to proving that there exist $u \in S_x$ such that $f(u) = -u$ because in that case, x, y and θ_u are on the same geodesic α_{x,θ_u}.

The following properties of f are obvious.

(i) For every $u \in S_x$, $f(u) \neq u$.

(ii) $f \circ f = Id$.

So f is an involution of the sphere without fixed point and for any such map, we claim that there exist u in the sphere such that $f(u) = -u$. In order to prove the claim, we follow a very similar argument in [16], theorem 1. We argue by contradiction. Let us assume that for every $u \in S_x$, $f(u) \neq -u$. The map $g : S_x \to S_x$ defined by $g(u) = \frac{f(u)+u}{\|f(u)+u\|}$, is then well defined and continuous. Let us remark that as for every $u \in S_x$, $f(u) \neq -u$, then f is homotopic to the Identity, and so is g. Moreover by (ii) we clearly have $g \circ f = g$, thus the map g factorizes through S_x/G_f where G_f is the group generated by the involution f. By (i) f has no fixed point thus S_x/G_f is a manifold and the projection $p : S_x \to S_x/G_f$ is a degre 2 map. Therefore, the induced endomorphism g_* on $H_{n-1}(S_x, \mathbb{Z}_2)$ is trivial, which contradicts the fact that g is homotopic to the Identity. □

Corollary 5.30. *Let $\mathcal{H}^{n-1} \subset \tilde{X}$ be an hypersurface with possibly non empty boundary $\partial\mathcal{H}^{n-1}$, such that for any $x \in \mathcal{H}^{n-1}$, $Jac_{n-1}\tilde{F}'(x) = Jac_{E_x}\tilde{F}'(x) = 1$, where E_x is the tangent space of $\mathcal{H}^{n-1}$ at x. Let us consider $x \in \mathcal{H}^{n-1}$ such that $dist_{\tilde{X}}(x, \partial\mathcal{H}^{n-1}) = r > 0$. Then, for any $x' \in \mathcal{H}^{n-1}$ with $dist_{\tilde{X}}(x, x') < r$, the geodesic $\alpha_{x,x'}$ joining x and x' is contained in $\mathcal{H}^{n-1}$. In particular, $\mathcal{H}^{n-1}$ is locally convex.*

Proof of the corollary: Let us fix $\theta \in \Lambda_{C'}$ and consider the vector field $\nabla B(y, \theta)$ in $\tilde{X}$. Let $x \in \mathcal{H}^{n-1}$. As $Jac_{E_x}\tilde{F}'(x) = 1$, we have by step 3 $\Lambda_{C'} = E_x(\infty)$. Then, for any $x \in \mathcal{H}^{n-1}$, $\nabla B(x, \theta)$ is tangent to $\mathcal{H}^{n-1}$, therefore the geodesic $\alpha_{x,\theta}$ satisfies $\alpha_{x,\theta}(t) \in \mathcal{H}^{n-1}$ for all $t \in [0, r)$. Let $x' \in \mathcal{H}^{n-1}$. By lemma 5.29, $\dot{\alpha}_{x,x'}(0) \in E_x$, therefore $\alpha_{x,x'} = \alpha_{x,\theta}$ and $\alpha_{x,x'}(t) \in \mathcal{H}^{n-1}$ for all $t \in [0, r)$. □

We now prove that Z_∞ is a totally geodesic hypersurface in $\tilde{X}/C'$. Let us recall that Z_∞ is a manifold which is smooth except at a singular subset of codimension at least 7. Let us consider a lift $\tilde{Z}_\infty \subset \tilde{X}$ of Z_∞ and denote $\tilde{Z}_\infty^{reg}$ (resp. $\tilde{Z}_\infty^{sing}$) the set of regular (resp.) singular points of $\tilde{Z}_\infty$.

Lemma 5.31. *$\tilde{Z}_\infty$ is a totally geodesic hypersurface in $\tilde{X}$.*

Proof: Let us consider a regular point $x \in \tilde{Z}_\infty^{reg}$. We shall show that for every point $x' \in \tilde{Z}_\infty^{reg}$ the geodesic segment joining x and x' is contained in $\tilde{Z}_\infty$, and as the set of regular points is dense in $\tilde{Z}_\infty$ (as the complementary of a subset of codimension at least 8), this will show that $\tilde{Z}_\infty$ is totally geodesic.

We claim that there exist a sequence $y_k \in \tilde{Z}_\infty^{reg}$ such that $\lim_{k\to\infty} y_k = x'$ and the geodesic segment joining x and y_k is contained in $\tilde{Z}_\infty$.

The claim immediately implies that the geodesic segment joining x and x' is contained in $\tilde{Z}_\infty$.

Let us prove the claim.

For $y \in \tilde{Z}_\infty^{reg}$ we consider $\alpha_{x,y}$ the geodesic joining x and y and define

$$t_y = inf\{t > 0, \alpha_{x,y}(t) \notin \tilde{Z}_\infty\} \tag{5.62}$$

As x is a regular point, by corollary 5.30, there exist $\epsilon > 0$ such that $t_y > \epsilon$.

In order to prove the claim, we argue by contradiction. Let us assume that there exist $r > 0$ such that for any $y \in B_{\tilde{X}}(x', r) \cap \tilde{Z}_\infty^{reg}$, $t_y < dist(x, y)$. By corollary 5.30 applied to $\tilde{Z}_\infty^{reg}$, we have $\alpha_{x,y}(t_y) \in \tilde{Z}_\infty^{sing}$. As the set of regular points is an open subset of $\tilde{Z}_\infty$, if r is small enough we have $B_{\tilde{X}}(x', r) \cap \tilde{Z}_\infty^{reg} = B_{\tilde{X}}(x', r) \cap \tilde{Z}_\infty$. We choose such an r and we consider the set S of all singular points contained in the union of all geodesic segments joining x to a point $y \in B_{\tilde{X}}(x', r) \cap Z_\infty$. Let us consider the map defined on S by

$$p(y) = \alpha_{x,y}(\epsilon).$$

As we already saw, for any $y \in B_{\tilde{X}}(x', r) \cap \tilde{Z}_\infty$, we have $t_y > \epsilon$, therefore the map p is distance decreasing and by assumption p is surjective onto an open subset of the sphere and $p(S)$ is homeomorphic to an open subset of $\mathbb{R}^{n-1}$, therefore the Hausdorff dimension of S is greater than or equal to $n-1$, which contradicts the fact that the singular set has codimension at least 8 in $\tilde{Z}_\infty$. □

The totally geodesic hypersurface $\tilde{Z}_\infty \subset \tilde{X}$ is preserved by C, and $\tilde{Z}_\infty / C$ is of minimal volume in its homology class.

Let us prove that $\tilde{Z}_\infty$ is isometric to the hyperbolic space $\mathbb{H}_{\mathbb{R}}^{n-1}$.

Lemma 5.32. *$\tilde{Z}_\infty$ is isometric to the hyperbolic space $\mathbb{H}_{\mathbb{R}}^{n-1}$.*

Proof :

As $\tilde{Z}_\infty / C$ is of minimal volume in its homology class, we have by Proposition 5.1, for all $x \in \tilde{Z}_\infty$, $Jac_{E_x}\tilde{F}(x) = 1$ and $\tilde{F}(x) = x$, where E_x is the tangent space of $\tilde{Z}_\infty$ at x. Moreover, we saw in the proof of proposition 5.1 that

$$H = \frac{1}{n-1} Id_{D\tilde{F}(x)(E_x)} = \frac{1}{n-1} Id_{E_x},$$

therefore we get from (4.11) and $\tilde{F}(x) = x$, that for all $u, v \in T_x\tilde{Z}_\infty$,

$$\int_{\partial\tilde{X}} [DdB_{(x,\theta)}(u, v) + DB_{(x,\theta)}(u)DB_{(x,\theta)}(v)] d\nu_x(\theta)$$

$$= \tilde{g}(u, v) \tag{5.63}$$

where $\tilde{g}$ is the metric on $\tilde{X}$. As $\tilde{Z}_\infty$ is totally geodesic, the relation (5.62) remains true with the Busemann function $B^{\tilde{Z}_\infty}$ of $\tilde{Z}_\infty$ instead of the Busemann function B of $\tilde{X}$:

$$\int_{\partial\tilde{X}} [DdB^{\tilde{Z}_\infty}_{(x,\theta)}(u,v) + DB^{\tilde{Z}_\infty}_{(x,\theta)}(u)DB^{\tilde{Z}_\infty}_{(x,\theta)}(v)$$

$$= \tilde{g}(u,v). \tag{5.64}$$

On the other hand, as $\tilde{Z}_\infty$ is totally geodesic, its sectional curvature is less than or equal to -1, thus by Rauch comparison theorem, we have

$$DdB^{\tilde{Z}_\infty}(x,\theta) + DB^{\tilde{Z}_\infty}(x,\theta) \otimes DB^{\tilde{Z}_\infty}(x,\theta) \geq \tilde{g}_{\tilde{Z}_\infty} \tag{5.65}$$

for all $\theta \in \partial\tilde{Z}_\infty = \Lambda_C$, where $\tilde{g}_{\tilde{Z}_\infty}$ is the restriction of $\tilde{g}$ to $\tilde{Z}_\infty$.

As the support of the measure ν_x is $\partial\tilde{Z}_\infty = \Lambda_C$ (by convex cocompactness of C) and the Busemann function is continuous, we get from (5.64) and (5.65) that for all $x \in \tilde{Z}_\infty$ and all $\theta \in \tilde{Z}_\infty$

$$DdB^{\tilde{Z}_\infty}(x,\theta) + DB^{\tilde{Z}_\infty}(x,\theta) \otimes DB^{\tilde{Z}_\infty}(x,\theta)$$

$$= \tilde{g}_{\tilde{Z}_\infty}(x). \tag{5.66}$$

and this last relation is characteristic of the hyperbolic space. $\square$

Step 6: Conclusion

So far we have shown that C preserves a totally geodesic copy of the hyperbolic space $\mathbb{H}^{n-1}_{\mathbb{R}} \subset \tilde{X}$ such that $\mathbb{H}^{n-1}_{\mathbb{R}}/C$ is compact.

Our goal now is to show that $Y =: \mathbb{H}^{n-1}_{\mathbb{R}}/C$ injects diffeomorphically in $X = \tilde{X}/\Gamma$ and separates X in two connected components R and S such that $A = \pi_1(R)$ and $B = \pi_1(S)$.

In order to do this, we will consider the Γ orbit of $\mathbb{H}^{n-1}_{\mathbb{R}}$ in $\tilde{X}$ and the two connected components U and V of $\tilde{X} - \Gamma\mathbb{H}^{n-1}_{\mathbb{R}}$ which are adjacent to $\mathbb{H}^{n-1}_{\mathbb{R}}$. The stabilizers $\bar{A}$, $\bar{B}$ and $\bar{C}$ of U, V and $\mathbb{H}^{n-1}_{\mathbb{R}}$ contain respectively A, B and C and the hypersurface $\mathbb{H}^{n-1}_{\mathbb{R}}/C$ injects in $X = \tilde{X}/\Gamma$ and separates X in two connected components R and S such that $\pi_1(R) = \bar{A}$ and $\pi_1(S) = \bar{B}$. We then show that $\bar{C} = C$, $\bar{A} = A$ and $\bar{B} = B$.

Let $\bar{C}$ be the stabilizer of $\mathbb{H}^{n-1}$, namely $\bar{C} = \{\gamma \in \Gamma, \gamma\mathbb{H}^{n-1} = \mathbb{H}^{n-1}\}$. We have $C \subset \bar{C}$ and as $\mathbb{H}^{n-1}/C$ is compact, so is $\mathbb{H}^{n-1}/\bar{C}$ and thus $[\bar{C} : C] < \infty$.

Let $p : \tilde{X}/C \to X = \tilde{X}/\Gamma$ and $\bar{p} : \tilde{X}/\bar{C} \to X = \tilde{X}/\Gamma$ the natural projections. We now show that the restriction of p to $\mathbb{H}^{n-1}/C$ is an embedding, thus $Y := p(\mathbb{H}^{n-1}/C)$ is a compact totally geodesic hypersurface of X.

In the section 2, we constructed a C-invariant hypersurface $\tilde{Z} \subset \tilde{X}$ such that $Z = \tilde{Z}/C \subset \tilde{X}/C$ is compact. The hypersurface is defined as $\tilde{Z} = \tilde{f}^{-1}(t_0)$ where $\tilde{f} : \tilde{X} \to T$ is an equivariant map onto the Bass-Serre tree associated to the amalgamation $A *_C B$ and t_0 belongs to that edge of T which is fixed by C.

Let us first show two lemmas.

Lemma 5.33. *The restriction of p to $\tilde{Z}/C$ is an embedding into $X = \tilde{X}/\Gamma$.*

Proof : Let $\gamma \in \Gamma$, z, z' in $\tilde{Z}$ such that $z' = \gamma z$. By equivariance,

$$\tilde{f}(\gamma z) = \gamma \tilde{f}(z) = \gamma t_0 = \tilde{f}(z') = t_0,$$

thus $\gamma \in C$.□

Lemma 5.34. *The restriction of $\bar{p}$ to $\mathbb{H}^{n-1}/\bar{C}$ is an embedding into $X = \tilde{X}/\Gamma$.*

Proof : Let us assume that there is a $\gamma \in \Gamma - \bar{C}$ such that $\gamma\mathbb{H}^{n-1} \cap \mathbb{H}^{n-1} \neq \emptyset$ and choose an $x \in \gamma\mathbb{H}^{n-1} \cap \mathbb{H}^{n-1}$. As $\gamma \notin \bar{C}$, there exist $u \in T_x\gamma\mathbb{H}^{n-1} - T_x\mathbb{H}^{n-1}$. We consider c_u the geodesic ray such that $\dot{c}_u(0) = u$. We know that $\tilde{Z}$ is contained in an ϵ-neighbourhood $\mathcal{U}_\epsilon\mathbb{H}^{n-1}$ of $\mathbb{H}^{n-1}$. The ϵ-neighbourhood $\mathcal{U}_\epsilon\mathbb{H}^{n-1}$ of $\mathbb{H}^{n-1}$ separates $\tilde{X}$ in two connected components U and V and for $t > 0$ large enough, we have, say, $c_u(t) \in U$ and $c_u(-t) \in V$.

Let $\tilde{Z}'$ be the connected component of $\tilde{Z}$ that we constructed at the end of section 2, whose stabilizer (or an index two subgroup of it) C' is such that $\tilde{Z}'/C'$ separates $\tilde{X}/C'$ in two unbounded connected components U'/C' and V'/C' where U' and V' are the two connected components of $\tilde{X} - \tilde{Z}'$.

We claim that $U \subset U'$ and $V \subset V'$ or the other way around. Indeed if not, U and V would be both contained in, say, U'. But in that case, V' would be contained in $\mathcal{U}_\epsilon H^{n-1}$ and therefore V'/C' would be bounded, which is a contradiction.

As $\gamma\tilde{Z}$ lies in the ϵ neighborhood of $\gamma\mathbb{H}^{n-1}$, there exist sequences z_k, z'_k in $\gamma\tilde{Z}$ such that $dist(z_k, c_u(k)) \leq \epsilon$ and $dist(z'_k, c_u(-k)) \leq \epsilon$. By proposition 5.13 and lemma 5.23, C' also acts cocompactly on $\mathbb{H}^{n-1}$, thus C' is of finite index in C, and therefore there are finitely many connected components of $\tilde{Z}$ and the same holds for $\gamma\tilde{Z}$. We thus can assume that the z_k's and z'_k's belong to a single connected component of $\gamma\tilde{Z}$. Let us consider a continuous path $\alpha \subset \gamma\tilde{Z}$ joining z_k and z'_k.

By construction the distance between $c_u(k)$ [resp. $c_u(-k)$] and $\mathbb{H}^{n-1}$ tends to infinity and thus, for k large enough, $z_k \in U$ and $z'_k \in V$ or the other way around. By the claim, we then have $z_k \in U'$ and $z'_k \in V'$, therefore the path α has to cross $\tilde{Z}'$ which contradicts the lemma (5.29) and ends the proof of the lemma 5.30. □

As we already saw, $\tilde{Z}$ has finitely many connected components, and so does $\tilde{X} - \tilde{Z}$. Let us write $\{W_j\}_{j=1,..,m}$ the connected components of $\tilde{X} - \tilde{Z}$. As C acts cocompactly on $\tilde{Z}$ and $\mathbb{H}^{n-1}$ there exist $\epsilon > 0$ such that $\mathbb{H}^{n-1} \subset$

$\mathcal{U}_\epsilon \tilde{Z}$ and $\tilde{Z} \subset \mathcal{U}_\epsilon \mathbb{H}^{n-1}$. Moreover $\mathcal{U}_\epsilon \mathbb{H}^{n-1}$ separates $\tilde{X}$ in two connected components U and V.

Lemma 5.35. *Let us consider ϵ such that $\tilde{Z} \subset \mathcal{U}_\epsilon \mathbb{H}^{n-1}$ and U and V the two connected components of $\tilde{X} - \mathcal{U}_\epsilon \mathbb{H}^{n-1}$. There are two distinct connected components W_1 and W_2 of $\tilde{X} - \tilde{Z}$ such that $U \subset W_1$ and $V \subset W_2$. Moreover, $\tilde{f}(W_1) \subset \tilde{T}_1$ and $\tilde{f}(W_2) \subset \tilde{T}_2$, where $\tilde{T}_1$ and $\tilde{T}_2$ are the two connected components of $\tilde{T} - \{t_0\}$.*

Proof : We argue by contradiction. Let us assume that U and V are contained in the same connected component W_1 of $\tilde{X} - \tilde{Z}$. Then, all other components W_j, $j \neq 1$, satisfy $W_j \subset \mathcal{U}_\epsilon \mathbb{H}^{n-1} \subset \mathcal{U}_{2\epsilon} \tilde{Z}$. Therefore, as C acts cocompactly on $\mathcal{U}_{2\epsilon} \tilde{Z}$, there exist a constant D such that for any $j \neq 1$, $max_{w \in W_j} dist_{\tilde{T}}(\tilde{f}(w), t_0) \leq D$. Thus, $\tilde{f}(W_1)$ is contained in one connected component of $\tilde{T} - \{t_0\}$ and $\tilde{f}(\cup_{j \neq 1} W_j)$, contained in the ball $B_{\tilde{T}}(t_0, D)$ of $\tilde{T}$ of radius D centered at t_0, is bounded. This is clearly impossible because $\tilde{T} - \{t_0\}$ has two unbounded connected components and $\tilde{f}$ is onto. □

Let us denote $\mathcal{A} = A\mathbb{H}^{n-1}$ the A-orbit of the C-invariant totally geodesic copy of the real hyperbolic space $\mathbb{H}^{n-1}$, and $\bar{A}$ the stabilizer of $\mathcal{A}$, ie. $\bar{A} = \{\gamma \in \Gamma, \gamma\mathcal{A} = \mathcal{A}\}$. We define in a similar way $\mathcal{B} = B\mathbb{H}^{n-1}$ and $\bar{B} = \{\gamma \in \Gamma, \gamma\mathcal{B} = \mathcal{B}\}$.

Let us recall that $\bar{C}$ is the stabilizer of $\mathbb{H}^{n-1}$ in Γ. We now prove the following

Lemma 5.36. *We have $\bar{A} = A\bar{C}$ and $\bar{B} = B\bar{C}$. Moreover, $\bar{A}$ and $\bar{B}$ are charactrized by $\bar{A} = \{\gamma \in \Gamma, \gamma\mathbb{H}^{n-1} \in \mathcal{A}\}$ and $\bar{B} = \{\gamma \in \Gamma, \gamma\mathbb{H}^{n-1} \in \mathcal{B}\}$.*

Proof : Let $\gamma' \in \bar{A}$, then $\gamma'\mathbb{H}^{n-1} \in \mathcal{A}$ and thus there exist $\gamma \in A$ such that $\gamma'\mathbb{H}^{n-1} = \gamma\mathbb{H}^{n-1}$, therefore $\gamma^{-1}\gamma' \in \bar{C}$, which proves the first part of the lemma.

Let us prove the second part of the lemma.

Let $\gamma' \in \Gamma$ be such that $\gamma'\mathbb{H}^{n-1} \in \mathcal{A}$. Then there exist $\gamma \in A$ such that $\gamma'\mathbb{H}^{n-1} = \gamma\mathbb{H}^{n-1}$, thus $\gamma^{-1}\gamma' \in \bar{C}$ and therefore $\gamma' \in \gamma\bar{C} \subset A\bar{C} = \bar{A}$. This proves one inclusion, the other inclusion being obvious. □

For each $\gamma \in \Gamma$, $\gamma\mathbb{H}^{n-1}$ separates $\tilde{X}$ in two connected components U_γ and V_γ.

Let us now prove the following lemma.

Lemma 5.37. *(i) Let $\gamma \in A$, [resp. $\gamma \in B$]. Then, we have $\mathcal{A} - \{\gamma\mathbb{H}^{n-1}\} \subset U_\gamma$ or $\mathcal{A} - \{\gamma\mathbb{H}^{n-1}\} \subset V_\gamma$, [resp. $\mathcal{B} - \{\gamma\mathbb{H}^{n-1}\} \subset U_\gamma$ or $\mathcal{B} - \{\gamma\mathbb{H}^{n-1}\} \subset V_\gamma$.]*

(ii) Let γ be an element of $\Gamma - \bar{A}$, [resp. $\Gamma - \bar{B}$]. Then $\mathcal{A} \subset U_\gamma$ or $\mathcal{A} \subset V_\gamma$, [resp. $\mathcal{B} \subset U_\gamma$ or $\mathcal{B} \subset V_\gamma$].

Proof : (i) We argue by contradiction. Let us consider $\gamma\mathbb{H}^{n-1}$, $\gamma'\mathbb{H}^{n-1}$ and $\gamma''\mathbb{H}^{n-1}$ three distinct elements in $\mathcal{A}$ such that $\gamma'\mathbb{H}^{n-1} \subset U_\gamma$ and $\gamma''\mathbb{H}^{n-1} \subset V_\gamma$. By equivariance we can assume γ is the identity. Let us recall that U and V are the two connected components of $\tilde{X} - \mathcal{U}_\epsilon \mathbb{H}^{n-1}$.

We then have $\gamma'\mathbb{H}^{n-1} \cap U \neq \emptyset$ and $\gamma''\mathbb{H}^{n-1} \cap V \neq \emptyset$, which implies $\gamma'\tilde{Z} \cap U \neq \emptyset$ and $\gamma''\tilde{Z} \cap V \neq \emptyset$.

By lemma (5.31), $U \subset W_1$ and $V \subset W_2$ where W_1 and W_2 are two connected components of $\tilde{X} - \tilde{Z}$ and $\tilde{f}(U) \subset \tilde{T}_1$ and $\tilde{f}(V) \subset \tilde{T}_2$, therefore, $\tilde{f}(U)$ contains $\gamma' t_0 \in \tilde{T}_1$ and $\tilde{f}(V)$ contains $\gamma'' t_0 \in \tilde{T}_2$. This is impossible because for all elements γ' and γ'' in A, $\gamma' t_0$ and $\gamma'' t_0$ belong to the same connected component of $\tilde{T} - \{t_0\}$.

(ii) Let us consider $\gamma \in \Gamma - \bar{A}$. We argue by contradiction. Let us assume there exist γ, γ' in A such that

$$\gamma'\mathbb{H}^{n-1} \subset U_\gamma$$

$$\gamma''\mathbb{H}^{n-1} \subset V_\gamma \tag{5.67}$$

Let $\epsilon > 0$ such that $\tilde{Z} \subset \mathcal{U}_\epsilon \mathbb{H}^{n-1}$ and U and V the connected component of $\tilde{X} - \mathcal{U}_\epsilon \mathbb{H}^{n-1}$. By lemma (5.31) we have $\tilde{f}(\gamma U) \subset \gamma \tilde{T}_1$ and $\tilde{f}(\gamma V) \subset \gamma \tilde{T}_2$, where $\gamma \tilde{T}_1$ and $\gamma \tilde{T}_2$ are the two connected components of $\tilde{T} - \{\gamma t_0\}$. By assumption (5.62), we have

$\gamma'\mathbb{H}^{n-1} \cap \gamma U \neq \emptyset$ and $\gamma''\mathbb{H}^{n-1} \cap \gamma V \neq \emptyset$, which implies $\gamma'\tilde{Z} \cap \gamma U \neq \emptyset$ and $\gamma''\tilde{Z} \cap \gamma V \neq \emptyset$, therefore $\gamma' t_0 \in \gamma \tilde{T}_1$ and $\gamma'' t_0 \in \gamma \tilde{T}_2$, which is impossible because in the tree $\tilde{T}$, the points $\gamma' t_0$ and $\gamma'' t_0$ belong to two adjacent edges.

□

By lemma (5.33) (i), for every γ in $\bar{A}$, [resp. $\bar{B}$], we can define U_γ as the connected component of $\tilde{X} - \mathbb{H}^{n-1}$ which contains all $\gamma'\mathbb{H}^{n-1}$ for all γ' in $\bar{A}$, [resp. $\bar{B}$], and $\gamma'\mathbb{H}^{n-1} \neq \gamma\mathbb{H}^{n-1}$.

Let us define

$$U_A := \cap_{\gamma \in A} U_\gamma. \tag{5.68}$$

By definition, U_A [resp. U_B]is a convex set in $\tilde{X}$ whose boundary is the collection $\mathcal{A}$, [resp. $\mathcal{B}$], of $\gamma\mathbb{H}^{n-1}$, γ in $\bar{A}$ [resp. $\bar{B}$] and by lemma (5.33) (ii), U_A and U_B are two disjoint connected components of $\tilde{X} - \Gamma\mathbb{H}^{n-1}$.

In fact, U_A [resp. U_B], is the convex hull of $\mathcal{A}$, [resp. $\mathcal{B}$], and $\bar{A}$, [resp. $\bar{B}$], is the stabilizer of U_A, [resp. U_B].

Lemma 5.38. *The closures of U_A and U_B intersect along $\mathbb{H}^{n-1}$ and $\bar{A} \cap \bar{B} = \bar{C}$. Moreover, no element of Γ sends U_A on U_B nor the other way around.*

Proof : The convex set U_A is the intersection of open half spaces U_γ, $\gamma \in \bar{A}$, and is delimited by the disjoint union of hyperplanes $\gamma\mathbb{H}^{n-1}$, for some $\gamma \in \bar{A}$. The same is true for U_B and as $U_A \cap U_B = \emptyset$, the closures of U_A and U_B can intersect only along one of the connected components of their boundaries, thus along $\mathbb{H}^{n-1}$ which is obviously in both closures. This proves the first part of the lemma, let us prove the second part. By lemma 5.32, $\bar{C} \subset \bar{A} \cap \bar{B}$. Conversely, let us take $\gamma \in \bar{A} \cap \bar{B}$, then γ preserves the closures of U_A and U_B, thus it preserves their intersection $\mathbb{H}^{n-1}$, and therefore $\gamma \in \bar{C}$.

Let us prove the last part of the lemma. Let γ be an element such that $\gamma U_A = U_B$. As $\mathbb{H}^{n-1}$ is one component of the boundary $\mathcal{B}$ of U_B, there exist one component $\gamma' \mathbb{H}^{n-1} \in \mathcal{A}$, γ' being in $\bar{A}$, such that $\gamma(\gamma'\mathbb{H}^{n-1}) = \mathbb{H}^{n-1}$. Therefore, $\gamma\gamma' \in \bar{C}$, thus $\gamma \in \bar{A}$. The same argument yields $\gamma^{-1} \in \bar{B}$, so $\gamma \in \bar{A} \cap \bar{B} = \bar{C}$ and γ preserves U_A and U_B, which contradicts our choice of γ.

□

The Γ-orbit of the closure of $U_A \cup U_B$ covers $\tilde{X}$. Let us construct a tree $\bar{T}$ embedded in $\tilde{X}$ in the following way: the set of vertices is the set of the connected components of $\tilde{X} - \Gamma\mathbb{H}^{n-1}$ and two vertices are joined by an edge if the boundaries of their corresponding connected components intersect non trivially in $\tilde{X}$. By construction Γ acts on $\bar{T}$, the stabilizers of the vertices a and b corresponding to U_A and U_B are $\bar{A}$ and $\bar{B}$, the stabilizer of the edge between a and b is $\bar{C}$ and a fundamental domain for this action is the segment joining a and b. By [14], I, 4, Theorem 6, the group Γ is the amalgamated product of $\bar{A}$ and $\bar{B}$ over $\bar{C}$.

We now claim that $\bar{A} = A$, $\bar{B} = B$ and $\bar{C} = C$.

As A, B and C are subgroups of $\bar{A}$, $\bar{B}$, and $\bar{C}$, the corresponding Mayer-Vietoris sequences of $A *_C B$ and $\bar{A} *_{\bar{C}} \bar{B}$ are related by the following commutative diagram

$$\begin{tikzcd}
H_n(A,\mathbb{R}) \oplus H_n(B,\mathbb{R}) \arrow[r] \arrow[d] & H_n(\Gamma,\mathbb{R}) \arrow[r] \arrow[d] & H_{n-1}(C,\mathbb{R}) \arrow[d] \\
H_n(\bar{A},\mathbb{R}) \oplus H_n(\bar{B},\mathbb{R}) \arrow[r] & H_n(\Gamma,\mathbb{R}) \arrow[r] & H_{n-1}(\bar{C},\mathbb{R}).
\end{tikzcd}$$

We know that the index $[\bar{C} : C]$ is finite, and by the lemma 5.32, the indices of A, B, and C in $\bar{A}$, $\bar{B}$ and $\bar{C}$ are finite and equal. On the other hand, the indices $[\Gamma : A]$ and $[\Gamma : B]$ are infinite by assumption, thus the previous diagram becomes

$$\begin{tikzcd}
0 \arrow[r] & H_n(\Gamma,\mathbb{R}) \arrow[r] \arrow[d] & H_{n-1}(C,\mathbb{R}) \arrow[d] \\
0 \arrow[r] & H_n(\Gamma,\mathbb{R}) \arrow[r] & H_{n-1}(\bar{C},\mathbb{R}).
\end{tikzcd}$$

Moreover, the map $H_n(\Gamma,\mathbb{R}) \to H_{n-1}(\bar{C},\mathbb{R})$ is bijective. Namely, the injectivity comes from the above diagram and the surjectivity from the fact that that the hypersurface $\mathbb{H}^{n-1}/\bar{C}$ bounds in $\mathbb{H}^n/A$ and $\mathbb{H}^n/B$ so that the map $H_{n-1}(C,\mathbb{R}) \to H_{n-1}(A,\mathbb{R}) \oplus H_{n-1}(B,\mathbb{R})$ is trivial. Therefore the index $[\bar{C} : C] = 1$ and we get $\bar{A} = A$, $\bar{B} = B$ and $\bar{C} = C$. □

6. Proof of the theorems 1.5 and 1.6.

The proof of theorem 1.5 is exactly the same as the proof of theorem 1.2. The actions of Γ on $\tilde{(X)}$ and T give rise to a continuous Γ-equivariant map $\tilde{f} : \tilde{X} \to T$. Like in section 2, we build an hypersurface $\tilde{f}^{-1}(t_0)$ where t_0 is a regular value of $\tilde{f}$ belonging the interior of an edge. As the edge separates the tree in two unbounded components, the section 2 applies and we get a subgroup C' of C, and an hypersurface $\tilde{Z}' \subset \tilde{X}/C'$ which is essential. Now, if the action of Γ is minimal, every edge separates T in two unbounded components. □

7. Appendix

The goal of this section is to give a proof of lemma 5.9. This lemma is contained in lemma 2.1, 5.1 and 5.2 of [4], but our situation being not exactly the same, we reproduce it down here for sake of completness.

Let us restate the lemma 5.9.

Lemma 7.1. *Let $\mathcal{L} \subset \partial\tilde{X}$ be a closed C'-invariant subset and $\theta_0 \in \mathcal{L}$. We assume that there exist a sequence of positive real numbers $\lambda_k \to \infty$ such that the sequence of pointed metric spaces $(\mathcal{L}, \lambda_k d, \theta_0)$ converges in the pointed Gromov-Hausdorff topology to $(S, \delta, 0)$ where $(S, \delta, 0)$ is a weak tangent of $(\partial\tilde{X}, d)$. We also assume that there exist positive constants C and δ, a sequence of points $\theta_0^k = \theta_0, \theta_1^k, \theta_2^k \in \mathcal{L}$ and a sequence of elements $\gamma_k \in C'$ such that $C^{-1}\lambda_k^{-1} \leq d(\theta_i^k, \theta_j^k) \leq C\lambda_k^{-1}$ and $d(\gamma_k\theta_i^k, \gamma_k\theta_j^k) \geq \delta$ for all $0 \leq i \neq j \leq 2$. Then, $\mathcal{L}$ is homeomorphic to the one point compactification $\hat{S}$ of S. In particular $\mathcal{L}$ is homeomorphic to $\partial\tilde{X}$.*

We first give a definition of pointed Hausdorff-Gromov convergence which is equivalent to the definition 5.7. We follow [4], paragraph 4.

A sequence of metric spaces (Z_k, d_k, z_k) converges to the metric space $(S, \delta, 0)$ if for every $R > 0$, and every $\epsilon > 0$, there exist an integer N, a subset $D \subset B_S(0, R)$, subsets $D_k \subset B_{Z_k}(z_k, R)$ and bijections $f_k : D_k \to D$ such that for $k \geq N$,

(i) $f_k(z_k) = 0$,

(ii) the set D is ϵ-dense in $B_S(0, R)$, and the sets D_k are ϵ-dense in $B_{Z_k}(z_k, R)$,

(iii) $|d_{Z_k}(x, y) - d_Z(f_k(x), f_k(y))| < \epsilon$,

where x, y belong to D_k.

Let us describe now the lemmas 2.1 and 5.1 following [4].

For a metric space (Z, d) the cross ratio of four points $\{z_i\}$, $i = 1, ...4$, is the quantity

$$[z_1, z_2, z_3, z_4] := \frac{d(z_1, z_3)d(z_2, z_4)}{d(z_1, z_4)d(z_2, z_3)} \tag{7.1}$$

Given two metric spaces X and Y, an homeomorphism $\eta : [0, \infty) \to [0, \infty)$, and an injective map $f : X \to Y$, we say that f is an η-quasi-Möbius map if for any four points $\{x_i\}$, $i = 1, .., 4$, in X, we have

$$[f(x_1), f(x_2), f(x_3), f(x_4)] \leq \eta([x_1, x_2, x_3, x_4]). \tag{7.2}$$

For example, any discrete cocompact group of isometries of $\tilde{X}$, where $\tilde{X}$ is a Cartan-Hadamard manifold with sectional curvature $K \leq -1$, is acting on the ideal boundary $(\partial\tilde{X}, d)$ endowed with the Gromov distance by η-quasi-Möbius transformations for some η.

Lemma 7.2 ([4], Lemma 2.1)**.** *Let (X, d_X) and (Y, d_Y) be two compact metric spaces, and for any integer k, $g_k : \tilde{D}_k \to Y$ an η-quasi-Möbius map defined on a subset $\tilde{D}_k$ of X. We assume that the Hausdorff distance between $\tilde{D}_k$ and X satisfies*

$$lim_{k\to\infty} dist_H(\tilde{D}_k, X) = 0$$

and that for any integer k, there exist points (x_1^k, x_2^k, x_3^k) in D_k and (y_1^k, y_2^k, y_3^k) in Y, such that $g_k(x_i^k) = y_i^k$ for $i \in \{1,2,3\}$, $d_X(x_i^k, x_j^k) \geq \delta$ and $d_Y(y_i^k, y_j^k) \geq \delta$ for $i, j \in \{1,2,3\}, i \neq j$, where δ is independant of k. Then a subsequence of g_k converges uniformly to a quasi-Möbius map $f : X \to Y$, ie. $lim_{k_j\to\infty} dist_H(g_{k_j}, f|_{\tilde{D}_{k_j}}) = 0$. If in addition, we suppose that

$$lim_{k\to\infty} dist_H(g_k(\tilde{D}_k), Y) = 0,$$

then the sequence $\{g_{k_j}\}$ converges uniformly to a quasi-Möbius homeomorphism $f : X \to Y$.

Before stating the second lemma, let us define a metric space Z to be uniformly perfect if there exist a constant $\lambda \geq 1$ such that for every $z \in Z$ and $0 < R < diam Z$, we have $\bar{B}(z, R) - B(z, \frac{R}{\lambda}) \neq \emptyset$.

Lemma 7.3 ([4], lemma 5.1)**.** *Let Z be a compact uniformly perfect metric space and G an η-quasi-Möbius action on Z. Suppose that for each integer k we are given a set D_k in a ball $B_k = B(z, R_k) \subset Z$ that is $(\epsilon_k R_k)$-dense in B_k, where $\epsilon_k > 0$, distinct points $x_1^k, x_2^k, x_3^k \in B(z, \lambda_k R_k)$, where $\lambda_k > 0$, with*

$$d_Z(x_i^k, x_j^k) \geq \delta_k R_k$$

for $i, j \in \{1,2,3\}, i \neq j$, where $\delta_k > 0$, and groups elements $\gamma_k \in G$ such that for $y_i^k := \gamma_k(x_i^k)$ we have,

$$d_Z(y_i^k, y_j^k) \geq \delta'$$

for $i, j \in \{1,2,3\}, i \neq j$, where δ' is independant of k. Let $D'_k = \gamma_k(D_k)$, and suppose that $\lambda_k \to 0$ when $k \to \infty$, and the sequence $\frac{\epsilon_k}{\delta_k^2}$ is bounded. Then $lim_{k\to\infty} dist_H(D'_k, Z) = 0$.

Let us go back to the proof of lemma 6.1. By definition of convergence, there exist a subsequence of $\{\lambda_k\}$, which we still denote by $\{\lambda_k\}$, subsets $\tilde{D}_k \subset B_S(0, k)$, $D_k \subset B_{\lambda_k \mathcal{L}}(\theta_0, k)$, where $\tilde{D}_k$ and D_k are minimal $1/k$-dense subsets of $B_S(0,k)$ and $B_{(\mathcal{L},\lambda_k d)}(\theta_0,k)$, and bijections $f_k : \tilde{D}_k \to D_k$ such that for all $x, y \in \tilde{D}_k$,

$$\frac{1}{2}\delta(x,y) \leq \lambda_k d(f_k(x), f_k(y)) \leq 2\delta(x,y), \tag{7.3}$$

cf. [4], (5.4).

We can suppose that the points $\theta_0^k := \theta_0$, θ_1^k, and θ_2^k in lemma 6.1 belong to the set D_k. By assumption there exist elements $\gamma_k \in C'$ and a constant δ such that

$$d(\gamma_k\theta_i^k, \gamma_k\theta_j^k) \geq \delta \tag{7.4}$$

for all $i, j \in \{0,1,2\}$.

The lemma 6.1 is a direct consequence of the lemma 6.2 applied to $(X, d_X) = (\hat{S}, \hat{\delta})$ and $(Y, d_Y) = (\mathcal{L}, d)$ and to the sequence of maps $g_k := \gamma_k \circ f_k$, where $\hat{S}$ is the one point compactification of S and $\hat{\delta}$ the distance on $\hat{S}$ associated to δ, cf. [4] Lemma 2.2.

Let us denote x_0^k, x_1^k, x_2^k be the points in S such that $f_k(x_i^k) = \theta_i^k$, for $i \in \{0,1,2\}$.

Let us check that the assumptions of lemma 6.2 are verified.

The fact that $lim_{k\to\infty} dist_H(\tilde{D}_k, \hat{S}) = 0$ comes the same way as in [4], (5.5).

By (6.3), we have, $\delta(x_i^k, x_j^k) \geq \frac{\lambda_k}{2} d(\theta_i^k, \theta_j^k)$ and by assumption we then get

$$\delta(x_i^k, x_j^k) \geq \frac{1}{2C}. \tag{7.5}$$

We then get the separation assumption on triples of points by choosing $\delta := inf\{D, \frac{1}{2C}\}$.

It remains to check the assumption on $g_k(\tilde{D}_k) = \gamma_k \circ f_k(\tilde{D}_k) = \gamma_k(D_k)$, namely,

$$lim_{k\to\infty} dist_H(\gamma_k(D_k), \Lambda_{C'}) = 0. \tag{7.6}$$

In order to prove the property (6.6), we want to apply the lemma 6.3, but as the set $(\mathcal{L}, d)$ is a priori not uniformly perfect, we shall replace the uniform perfectness by the fact that $(\mathcal{L}, \lambda_k d, \theta_0)$ converges to a space $(S, \delta_0, 0)$, which is uniformly perfect, cf. ().

We will show the

Lemma 7.4. *We consider the subsets $\tilde{D}_k \subset B_S(0,k)$ and $D_k \subset B_{\lambda_k\mathcal{L}}(\theta_0,k)$, where $\tilde{D}_k$ and D_k are $1/k$-dense subsets of $B_S(0,k)$ and $B_{(\mathcal{L},\lambda_k d)}(\theta_0,k)$, and the bijections $f_k : \tilde{D}_k \to D_k$ coming from the convergence of the sequence of pointed metric spaces $(\mathcal{L}, \lambda_k d, \theta_0)$ to $(S, \delta, 0)$ where $(S, \delta, 0)$ is a weak tangent of $(\partial\tilde{X}, d)$. We also assume that there exist positive constants C and δ, a* *sequence of points* $\theta_1^k, \theta_2^k \in \Lambda_{C'}$ *and a sequence of elements* $\gamma_k \in C'$ *such that* $C^{-1}\lambda_k^{-1} \leq d(\theta_i^k, \theta_j^k) \leq C\lambda_k^{-1}$ *and* $d(\gamma_k\theta_i^k, \gamma_k\theta_j^k) \geq \delta$ *for all* $0 \leq i \neq j \leq 2$. *Then, the Hausdorf distance* $dist_H(\gamma_k D_k, \mathcal{L})$ *tends to* 0 *as* k *tends to infinity.*

Proof : The proof is word by word the same as the proof of lemma 6.3, ie. lemma 5.1 (i) of [4] with a difference in case 2).

We have $B_{\lambda_k\mathcal{L}}(\theta_0, k) = B_{\mathcal{L}}(\theta_0, \frac{k}{\lambda_k})$ and $D_k \subset B_{\lambda_k\mathcal{L}}(\theta_0, k)$ an $\frac{1}{k}$-dense subset, for the metric $\lambda_k d$. In term of the distance d, the set D_k is $(\epsilon_k R_k)$-dense in $B_{\mathcal{L}}(\theta_0, R_k)$, where $R_k := \frac{k}{\lambda_k}$ and $\epsilon_k := \frac{1}{k^2}$. By assumption, the points $\theta_0^k = \theta_0, \theta_1^k, \theta_2^k$ belong to $B_{\mathcal{L}}(\theta_0, \mu_k R_k)$, and satisfy

$$d(\theta_i^k, \theta_j^k) \geq \delta_k R_k \tag{7.7}$$

where $\delta_k := \frac{1}{Ck}$, and $\mu_k := \frac{C}{k}$.

The points $\gamma_k\theta_i^k$ satisfy

$$d(\gamma_k\theta_i^k, \gamma_k\theta_j^k) \geq \delta, \tag{7.8}$$

and $\frac{\epsilon_k}{\delta_k^2} = C^2$ is bounded.

Let us consider a point $\theta \in \mathcal{L}$. We want to approximate it by a point of $\gamma_k D_k$.

We can write $\theta = \gamma_k\theta_k$, for some $\theta_k \in \Lambda_{C'}$. There are two cases.

Case 1). For infinitely many indices k, $\theta_k \in B_{\mathcal{L}}(\theta_0, R_k)$. We work in that case for these indices k, thus there are points $\theta_k' \in D_k \cap B_{\mathcal{L}}(\theta_0, R_k)$, with $d(\theta_k, \theta_k') \leq \epsilon_k R_k$.

Since the distance between the θ_i^k's is bounded below by $\delta_k R_k$, we can find at least two of them which we call a_k and b_k, such that

$$d(\theta_k, b_k) \geq \frac{\delta_k R_k}{2}$$

and,

$$d(\theta_k', a_k) \geq \frac{\delta_k R_k}{2}. \tag{7.9}$$

As C' is contained in the cocompact group Γ, it acts in a quasi-Möbius way on $(\partial\tilde{X}, d)$

thus,

$$\frac{d(\gamma_k\theta_k', \gamma_k\theta_k)d(\gamma_k a_k, \gamma_k b_k)}{d(\gamma_k\theta_k', \gamma_k b_k)d(\gamma_k a_k, \gamma_k\theta_k)} \leq \eta(\frac{d(\theta_k', \theta_k)d(a_k, b_k)}{d(\theta_k', b_k)d(\theta_k, a_k)}) \tag{7.10}$$

for some homeomorphism $\eta : [0, \infty) \to [0, \infty)$. This implies

$$d(\gamma_k\theta_k', \gamma_k\theta_k) \leq \frac{(diam\mathcal{L})^2\eta(8\epsilon_k\mu_k/\delta_k^2)}{\delta}, \tag{7.11}$$

therefore $d(\gamma_k\theta_k', \gamma_k\theta_k)$ tends to zero as k tends to infinity.

Case 2) . For all but finitely many indices k, $\theta_k \notin B_{\mathcal{L}}(\theta_0, R_k)$.

We work with these indices k such that $\theta_k \notin B_{\mathcal{L}}(\theta_0, R_k)$.

We know that ϵ_k/δ_k^2 is bounded above independantly of k, and by assumption, $\delta_k \leq 2\mu_k$.

We claim that there exist $\xi_k \in D_k$ and a positive constant c_0 such that for all k,

$$\frac{d(\xi_k, \theta_0)}{R_k} \geq c_0 \tag{7.12}$$

let us prove the claim.

On one hand, as $(\partial\tilde{X}, d)$ is uniformly perfect, and so is it's weak tangent (S, δ) because the one point compactification $(\hat{S}, \hat{\delta})$ of (S, δ) is quasi-Möbius homeomorphic to $(\partial\tilde{X}, d)$, therefore there exist a constant $C_0 \in [0, 1)$ such that for every $x \in S$ and $0 < R < diam S$, we have

$$\bar{B}_{(S,\delta)}(0, R) - B_{(S,\delta)}(0, C_0 R) \neq \emptyset. \tag{7.13}$$

On the other hand, $(\mathcal{L}, \lambda_k d, \theta_0)$ converges to $(S, \delta, 0)$. After reindexing the sequence $\{\lambda_k\}$, we have for each $\epsilon > 0$ a map $g_k : B_{\lambda_k \mathcal{L}}(\theta_0, k) \to S$ such that
(i) $g_k(\theta_0) = 0$,
for any two points θ and θ' in $B_{\lambda_k \mathcal{L}}(\theta_0, k)$,
(ii) $|\delta(g_k(\theta), g_k(\theta')) - \lambda_k d(\theta, \theta')| \leq \epsilon$,
(iii) the ϵ-neighborhood of $g_k(B_{\lambda_k \mathcal{L}}(\theta_0, k))$ contains $B_{(S,\delta)}(0, k - \epsilon)$.

By (iii), we have

$$\bar{B}_{(S,\delta)}(0, k - \epsilon) \subset \mathcal{U}_\epsilon^{(S,\delta)} g_k(\bar{B}_{\lambda_k \mathcal{L}}(\theta_0, k)). \tag{7.14}$$

By (6.13) there exist $y_k \in \bar{B}_{(S,\delta)}(0, k - \epsilon) - B_{(S,\delta)}(0, C_0(k - \epsilon))$, and by (6.14) there exist $\xi_k' \in \bar{B}_{\lambda_k \mathcal{L}}(\theta_0, k)$ such that

$$\delta(y_k, g_k(\xi_k')) \leq \epsilon. \tag{7.15}$$

We now evaluate $d(y_k, g_k(\xi_k'))$. By the above properties (i), (ii), (6.15) and the triangle inequality we have

$$\lambda_k d(\xi_k', \theta_0) \geq \delta(g_k(\xi_k'), 0) - \epsilon \geq \delta(y_k, 0) - \delta(y_k, g_k(\xi_k')) - \epsilon$$

$$\geq C_0(k - \epsilon) - 2\epsilon. \tag{7.16}$$

As D_k is $\epsilon_k R_k$-dense in $B_{(\mathcal{L},d)}(\theta_0, k/\lambda_k)$, there exist $\xi_k \in D_k$ such that $d(\xi_k, \xi_k') \leq \epsilon_k R_k = \frac{k\epsilon_k}{\lambda_k}$.

Let us denote $c_0 = C_0/2$. For k large enough we have $\frac{C_0(k-\epsilon)-2\epsilon-k\epsilon_k}{\lambda_k} \geq \frac{c_0 k}{\lambda_k}$, therefore by (6.16) we get

$$d(\xi_k, \theta_0) \geq d(\xi'_k, \theta_0) - d(\xi'_k, \xi_k) \geq c_0 R_k, \tag{7.17}$$

which proves the claim.

We can assume that for k large enough, $\mu_k < c_0/2 < 1/2$.

We choose $a_k = \theta_1^k$ and $b_k = \theta_2^k$, and we get

$$\frac{d(\gamma_k\xi_k, \gamma_k\theta_k)d(\gamma_k a_k, \gamma_k b_k)}{d(\gamma_k\xi_k, \gamma_k b_k)d(\gamma_k a_k, \gamma_k\theta_k)} \leq \eta(\frac{d(\xi_k, \theta_k)d(a_k, b_k)}{d(\xi_k, b_k)d(\theta_k, a_k)})$$

$$\leq \eta(\frac{4\mu_k d(\theta_k, \theta_0^k)}{(d(\theta_k, \theta_0^k) - \mu_k R_k))(c_0 - \mu_k)})$$

$$\leq \eta(16\mu_k/c_0). \tag{7.18}$$

We get

$$d(\gamma_k\theta_k, \gamma_k\xi_k) \leq (diam\Lambda_{C'})^2 \eta(16\mu_k/c_0)/\delta.$$

□

References

[1] G. Besson, G. Courtois, S. Gallot, Lemme de Schwarz réel et applications géométriques, Acta Mathematica, 183, (1999), 145-169.

[2] G. Besson, G. Courtois, S. Gallot, Hyperbolic manifolds, amalgamated products and crtical exponents., C.R.MathAcad.Sci.Paris, 336, (2003), no. 3, 257-261.

[3] G. Besson, G. Courtois, S. Gallot,Entropies et rigidités des espaces localement symétriques de courbure négative, Geom. Funct. Anal. 5 (1995),731-799.

[4] M. Bonk, B. Kleiner, Rigidity for quasi-Möbius group actions, J. of Diff. Geometry 61, vol. 1, (2002), 81-106.

[5] R. Bott, L.W. Tu, Differential forms in algebraic topology, Springer-Verlag, New-York, (1982).

[6] K. S. Brown, Cohomology of groups, Springer-Verlag, (1982).

[7] D. Burago, Y. Burago, S. Ivanov, A course in Metric Geometry, Graduate Studies in Mathematics, vol. 33, American Mathematical Society, (2001).

[8] G. Carron, E. Pedon, On the differential form spectrum of hyperbolic manifold, to appear.

[9] M. Gromov, Filling riemannian manifolds, J. of Diff. Geometry 18, (1983), 1-147.

[10] V. Kaimanovich, Invariant measures of the geodesic flow and measure at infinity of negatively curved manifolds, Ann. Inst. H.Poincaré Phys. Theor. 53, 4, 361-393, (1994).

[11] A. Lubotsky, Free quotients and the first Betti number of some hyperbolic manifolds, Transformation groups vol. 1 (no 1, 2), (1996), 71-82.

[12] F. Morgan, Geometric measure theory, Academic Press, 1988.

[13] T.Roblin, Ergodicité et équidistribution en courbure négative, Mémoires de la Soc. Math. de France 95, 2003.

[14] J. P. Serre, Arbres, amalgames, SL_2, Société mathématique de France, Astérisque 46, (1977).

[15] Y. Shalom, Rigidity, unitary representations of semisimple groups, and fundamental groups of manifolds with rank one transformation group, Annals of Mathematics 152, (2000), 113-182.

[16] Z.I. Szabó, A short topological proof for the symmetry of @ point homogeneous spaces, Invent. Math. 106, 1991, 61-64.

[17] D. Sullivan Related aspects of positivity in riemannian geometry, J. Differetial Geometry 25, 1987, 327-351.

[18] D. Sullivan, Entropy, hausdorff measure old and new and limit sets of geometrically finite Kleinian groups, Acta Math. 153 (1984) 259-277.

[19] J. Väisälä, Quasi-Möbius maps, Journal d'Analyse Mathématique, vol. 44, 218-234, (1984/85).

[20] C. Yue, Then ergodic theory of discrete isometry groups on manifolds of negative curvature, Trans. Amer. Math. Soc. 348, (1996), 4965-5005.

[21] A. Zeghib, Laminations et hypersurfaces géodésiques des variétés hyperboliques, Ann. Sci. Ecole Norm. Sup. (4), no 2, 1991, 171-188.

Gérard Besson, Institut Fourier, UMR 5582 CNRS, BP 74 38402 St Martin d'Hères, France
E-mail address: `G.Besson@ujf-grenoble.fr`

Gilles Courtois, CMLS, École Polytechnique, UMR 7640 CNRS, 91128 Palaiseau, France
E-mail address: `courtois@math.polytechnique.fr`

Sylvain Gallot, Institut Fourier, UMR 5582 CNRS, BP 74 38402 St Martin d'Hères, France
E-mail address: `sylvestre.gallot@ujf-grenoble.fr`